



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C07D 417/12, 339/04, C07C 327/18, 327/36, A61K 31/216, 31/192, 31/427, 31/385</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/53601</b> <b>(43) International Publication Date:</b> 14 September 2000 (14.09.00)
<b>(21) International Application Number:</b> PCT/US00/05895 <b>(22) International Filing Date:</b> 7 March 2000 (07.03.00)  <b>(30) Priority Data:</b> 09/264,370           8 March 1999 (08.03.99)       US 09/497,324           3 February 2000 (03.02.00)       US  <b>(71) Applicant (for all designated States except US):</b> THE UNIVERSITY OF MISSISSIPPI [US/US]; 1006 Natural Products Center, University, MS 38677 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> PERSHADSINGH, Harrihar, A. [US/US]; 404 Windsor Park Drive, Bakersfield, CA 93311 (US). AVERY, Mitchell, A. [US/US]; 303 Woodland Hills Drive, Oxford, MS 38655 (US).  <b>(74) Agents:</b> SNYDER, Joseph, R. et al.; Townsend and Townsend and Crew LLP, Two Embarcadero Center, 8th floor, San Francisco, CA 94111 (US).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> 1,2-DITHIOLANE DERIVATIVES  <b>(57) Abstract</b> <p>This invention provides new thiazolindinedione derivatives and new arylacetic acid derivatives. These compounds are useful for the treatment of cardiovascular diseases, certain endocrine diseases, certain inflammatory diseases, certain neoplastic (malignant) and non-malignant proliferative diseases, certain neuro-psychiatric disorders, certain viral diseases, and diseases associated with these viral infections as discussed herein.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

<b>AL</b>	Albania	<b>ES</b>	Spain	<b>LS</b>	Lesotho	<b>SI</b>	Slovenia
<b>AM</b>	Armenia	<b>FI</b>	Finland	<b>LT</b>	Lithuania	<b>SK</b>	Slovakia
<b>AT</b>	Austria	<b>FR</b>	France	<b>LU</b>	Luxembourg	<b>SN</b>	Senegal
<b>AU</b>	Australia	<b>GA</b>	Gabon	<b>LV</b>	Latvia	<b>SZ</b>	Swaziland
<b>AZ</b>	Azerbaijan	<b>GB</b>	United Kingdom	<b>MC</b>	Monaco	<b>TD</b>	Chad
<b>BA</b>	Bosnia and Herzegovina	<b>GE</b>	Georgia	<b>MD</b>	Republic of Moldova	<b>TG</b>	Togo
<b>BB</b>	Barbados	<b>GH</b>	Ghana	<b>MG</b>	Madagascar	<b>TJ</b>	Tajikistan
<b>BE</b>	Belgium	<b>GN</b>	Guinea	<b>MK</b>	The former Yugoslav Republic of Macedonia	<b>TM</b>	Turkmenistan
<b>BF</b>	Burkina Faso	<b>GR</b>	Greece			<b>TR</b>	Turkey
<b>BG</b>	Bulgaria	<b>HU</b>	Hungary	<b>ML</b>	Mali	<b>TT</b>	Trinidad and Tobago
<b>BJ</b>	Benin	<b>IE</b>	Ireland	<b>MN</b>	Mongolia	<b>UA</b>	Ukraine
<b>BR</b>	Brazil	<b>IL</b>	Israel	<b>MR</b>	Mauritania	<b>UG</b>	Uganda
<b>BY</b>	Belarus	<b>IS</b>	Iceland	<b>MW</b>	Malawi	<b>US</b>	United States of America
<b>CA</b>	Canada	<b>IT</b>	Italy	<b>MX</b>	Mexico	<b>UZ</b>	Uzbekistan
<b>CF</b>	Central African Republic	<b>JP</b>	Japan	<b>NE</b>	Niger	<b>VN</b>	Viet Nam
<b>CG</b>	Congo	<b>KE</b>	Kenya	<b>NL</b>	Netherlands	<b>YU</b>	Yugoslavia
<b>CH</b>	Switzerland	<b>KG</b>	Kyrgyzstan	<b>NO</b>	Norway	<b>ZW</b>	Zimbabwe
<b>CI</b>	Côte d'Ivoire	<b>KP</b>	Democratic People's Republic of Korea	<b>NZ</b>	New Zealand		
<b>CM</b>	Cameroon		Republic of Korea	<b>PL</b>	Poland		
<b>CN</b>	China	<b>KR</b>	Republic of Korea	<b>PT</b>	Portugal		
<b>CU</b>	Cuba	<b>KZ</b>	Kazakstan	<b>RO</b>	Romania		
<b>CZ</b>	Czech Republic	<b>LC</b>	Saint Lucia	<b>RU</b>	Russian Federation		
<b>DE</b>	Germany	<b>LI</b>	Liechtenstein	<b>SD</b>	Sudan		
<b>DK</b>	Denmark	<b>LK</b>	Sri Lanka	<b>SE</b>	Sweden		
<b>EE</b>	Estonia	<b>LR</b>	Liberia	<b>SG</b>	Singapore		

## 1,2-DITHIOLANE DERIVATIVES

### 5 CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims priority to U.S. Patent Application Serial No. 09/264,370, filed March 8, 1999, and U.S. Patent Application Serial No. 09/497,324, filed February 3, 2000, the disclosures of which are incorporated by reference in their entirety for all purposes.

### 10 BACKGROUND OF THE INVENTION

Peroxisome proliferator-activated receptors (PPARs) are members of the nuclear receptor superfamily of ligand-activated transcription factors. Three subtypes of PPARs have been cloned from the mouse and human: *i.e.*, PPAR $\alpha$ , PPAR $\gamma$  and PPAR $\delta$ . The PPARs are believed to play a role in the regulation of lipid metabolism. They can be  
15 activated by high concentration of fatty acids and have been shown to regulate the expression levels of fatty acid binding proteins or enzymes involved in fatty acid oxidation.

It has previously been discovered that a certain class of thiazolidinediones are selective PPAR $\gamma$  agonists (*see*, Willson *et. al.*, *J. Med. Chem.* (1996) 39:665-668).  
20 Thiazolidinediones are a class of oral insulin-sensitizing agents that improve glucose utilization without stimulating insulin release. For instance, U.S. Patent No. 4,287,200, discloses certain thiazolidine derivatives having the ability to lower blood glucose levels. In addition, U.S. Patent No. 4,572,912, discloses thiazolidinedione derivatives having the ability to lower blood lipid and blood glucose levels. These compounds were shown  
25 to have the ability to decrease the levels of blood lipid peroxides, blood triglycerides and blood cholesterol.

Moreover, U.S. Patent No. 5,338,855, discloses thiazolidine derivatives containing a quinone moiety. These compounds were shown to have the ability to reduce insulin resistance in the peripheral tissues and possess the ability to suppress hepatic  
30 gluconeogenesis in the liver.


In addition to being anti-diabetic agents which can lower the concentration of glucose and lipids in the blood, U.S. Patent No. 5,594,015 discloses thiazolidine derivatives as being effective in the treatment of hyperproliferation of epithelial cell conditions, such as psoriatic activity.

5 The anti-diabetic effect of the thiazolidinediones and their PPAR $\gamma$  agonist activity has implicated PPAR $\gamma$  as the molecular target for the anti-diabetic effects of thiazolidinediones. PPAR $\gamma$  is predominately expressed in adipose tissue and has been implicated as a master regulator of adipocyte differentiation in pre-adipose cell lines.

10 In view of the role PPAR $\gamma$  plays in regulation of lipid and glucose metabolism and the agonistic behavior of thiazolidinediones, there remains a need in the art for new thiazolidinedione derivatives and more effective therapies for diabetes and other ailments. The present invention fulfills these and other needs.

### SUMMARY OF THE INVENTION

15 This invention provides new thiazolidinedione derivatives. As such, in one aspect, the present invention provides compounds of Formula I:




In Formula I, R<sup>1</sup> and R<sup>2</sup> are each independently a functional group including, but not limited to, hydrogen, C(O)-R<sup>6</sup> and C(S)-R<sup>6</sup>. R<sup>6</sup>, in Formula I, is a functional group including, but not limited to, hydrogen, (C<sub>1</sub>-C<sub>12</sub>)alkyl, aryl, arylalkyl, (C<sub>1</sub>-C<sub>12</sub>)carboxyl, (C<sub>1</sub>-C<sub>12</sub>)NHR<sup>7</sup>, (C<sub>1</sub>-C<sub>12</sub>)NR<sup>7</sup>R<sup>8</sup>, OR<sup>7</sup>, NHR<sup>7</sup>, SR<sup>7</sup>, NR<sup>7</sup>R<sup>8</sup>. R<sup>7</sup> and R<sup>8</sup>, in Formula I, are each independently functional groups including, but not limited to, hydrogen, (C<sub>1</sub>-C<sub>12</sub>)alkyl, aryl and arylalkyl. The bond bisecting the functional group containing two sulfurs, indicates that the bond can be attached at anyone of the 3 carbon atoms between the two sulfur atoms.

In an alternative embodiment, R<sup>1</sup> and R<sup>2</sup> together with the sulfurs to which they are bound join to form a 1,2-dithiolane ring. In this embodiment, the 1,2-dithiolane ring can be substituted at the 3-position or the 4-position.

30 X, in Formula I, is a functional group including, but not limited to O, NR, C(O)O, OC(O)O and C(O)NR, wherein R is a functional group including, but not limited

to, hydrogen and optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl. Y, in Formula I, is a functional group including, but not limited to, O, S and NR<sup>6</sup>, wherein R<sup>6</sup> is a functional group including, but not limited to, hydrogen and optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl. In Formula I, the index "n" is an integer from 2 to 14; the index "m" is an integer from 0 to 14; the index "q" is an integer from 0 to 1; and the index "t" is an integer from 0 to 1, provided when m is 0 then q is 0, or a pharmaceutical acceptable salt or solvate thereof. Suitable salts include, but are not limited to, sodium, potassium and ammonium.

In another embodiment, the present invention provides a compound of Formula II:



In Formula II, R<sup>1</sup> and R<sup>2</sup> are each independently a functional group including, but not limited to, hydrogen, C(O)-R<sup>6</sup> and C(S)-R<sup>6</sup>. R<sup>6</sup>, in Formula II, is a functional group including, but not limited to, hydrogen, (C<sub>1</sub>-C<sub>12</sub>)alkyl, aryl, arylalkyl, (C<sub>1</sub>-C<sub>12</sub>)carboxyl, (C<sub>1</sub>-C<sub>12</sub>)NHR<sup>7</sup>, (C<sub>1</sub>-C<sub>12</sub>)NR<sup>7</sup>R<sup>8</sup>, OR<sup>7</sup>, NHR<sup>7</sup>, SR<sup>7</sup>, NR<sup>7</sup>R<sup>8</sup>. R<sup>7</sup> and R<sup>8</sup>, in Formula II, are each independently a functional group including, but not limited to, hydrogen, (C<sub>1</sub>-C<sub>12</sub>)alkyl, aryl and arylalkyl. The bond bisecting the functional group containing two sulfurs, indicates that the bond can be attached at any of the 3 carbon atoms between the two sulfur atoms.

In an alternative embodiment, R<sup>1</sup> and R<sup>2</sup> together with the sulfurs to which they are bound join to form a 1,2-dithiolane ring. The 1,2 dithiolane ring can be substituted at the 3-position or at the 4-position.

X, in Formula II, is functional group including, but not limited to O, NR, C(O)O, OC(O)O and C(O)NR, wherein R is a functional group including, but not limited to, hydrogen, optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl and optionally substituted aryl. Y, in Formula II, is a functional group including, but not limited to, O, S and NR<sup>6</sup>, wherein R<sup>6</sup> is a functional group including, but not limited to, hydrogen and optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl. R<sup>4</sup>, in Formula II, is a functional group including, but not limited to, hydrogen, halogen, optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl and optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkoxy. R<sup>5</sup>, in Formula II, is a functional group including, but not limited to, hydrogen and optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl. In Formula II, the index "n" is an integer from 2 to 14; the index "m" is an integer from 0 to 14; and the index "q" is an integer from 0 to 1,

provided when m is 0 then q is 0, or a pharmaceutical acceptable salt or solvate thereof. Suitable salts include, but are not limited to, sodium, potassium and ammonium.

In another aspect, the present invention relates to a pharmaceutical composition comprising a compound of the Formula I wherein R, R<sup>1</sup>, R<sup>2</sup>, X, Y, R<sup>3</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup>, n, m, q and t have the same meaning as defined above, or a pharmaceutically acceptable salt or solvate thereof, and a pharmaceutical acceptable carrier.

In another aspect, the present invention relates to a pharmaceutical composition comprising a compound of the Formula II wherein R, R<sup>1</sup>, R<sup>2</sup>, X, Y, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup>, n, m, and q have the same meaning as defined above, or a pharmaceutically acceptable salt or solvate thereof, and a pharmaceutical acceptable carrier.

In yet another aspect, the present invention relates to a method of treating a PPAR $\gamma$  mediated disease or oxidative stress, comprising administering a therapeutically effective amount of a compound of Formulae I, II or mixtures thereof, to an individual suffering from a PPAR $\gamma$  mediated disease. In other aspects, this invention provides methods for synthesizing the compounds of Formulae I and II.

### BRIEF DESCRIPTION OF THE DRAWINGS

**Figure 1** illustrates a method to synthesize compounds of Group I of this invention. The following reagents are used: a) BrCH<sub>2</sub>CH<sub>2</sub>Br, K<sub>2</sub>CO<sub>3</sub>, acetone; b) MeNH<sub>2</sub>, H<sub>2</sub>O, MeOH; c) thiazolidine-2,4-dione, piperidine, THF; d) H<sub>2</sub>, Pd/C, MeOH; e) R-(+)- $\alpha$ -lipoic acid, DCC, pyridine, CH<sub>2</sub>Cl<sub>2</sub>; f) NaBH<sub>4</sub>, MeOH.

**Figure 2A-C** illustrates intermediate compounds and synthesis methods of this invention. Panel A shows chemical intermediates; panel B shows chemical intermediates; and panel C shows a rutin reaction. The following reagents are used: a) ClCH<sub>2</sub>CH<sub>2</sub>OH, NaOH ; b) H<sub>3</sub>O<sup>+</sup>.

**Figure 3** illustrates a method to synthesize compounds of Group I of this invention. The following reagents are used: a) HOCH<sub>2</sub>CH<sub>2</sub>Cl, K<sub>2</sub>CO<sub>3</sub>, acetone; b) Thiazolidine-2,4-dione, piperidine, THF; c) H<sub>2</sub>, Pd/C, MeOH; d) R-(+)- $\alpha$ -Lipoic Acid, DCC, pyridine, CH<sub>2</sub>Cl<sub>2</sub>; e) NaBH<sub>4</sub>, MeOH.

**Figure 4** illustrates a method to synthesize compounds of Group I of this invention. The following reagents are used: a) LiBH<sub>4</sub>, THF; air oxidation; b) 2NaH,

COCl<sub>2</sub>, THF; c) thiazolidine-2,4-dione, piperidine, THF; d) H<sub>2</sub>, Pd/C, MeOH; e) Air oxidation.

**Figure 5** illustrates a method to synthesize compounds of Group I of this invention. The following reagents are used: a) LiBH<sub>4</sub>, THF; air oxidation; b) 2NaH, COCl<sub>2</sub>, THF; c) thiazolidine-2,4-dione, piperidine, THF; d) H<sub>2</sub>, Pd/C, MeOH; e) Air oxidation.

**Figure 6** illustrates a method to synthesize compounds of Group II of this invention. The following reagents are used: a) R-(+)-lipoic acid, DCC, pyridine; b) thiazolidine-2,4-dione, piperidine, THF; c) H<sub>2</sub>, Pd/C, MeOH; d) air oxidation.

**Figure 7** illustrates a method to synthesize compounds of Group II of this invention. The following reagents are used: a) LiBH<sub>4</sub>, THF; air oxidation; b) *p*-MePhSO<sub>2</sub>Cl, NaH; then NaH, 4-hydroxybenzaldehyde; c) thiazolidine-2,4-dione, piperidine, THF; d) H<sub>2</sub>, Pd/C, MeOH; e) air oxidation.

**Figure 8** illustrates a method to synthesize compounds of Group III of this invention. The following reagents are used: a) DCC, pyridine; then MeNH(CH<sub>2</sub>)<sub>3</sub>OH; b) NaH, 2-Fluoro-N-methylpyridinium iodide; then DMSO, Et<sub>3</sub>N; c) thiazolidine-2,4-dione, 2BuLi, THF; then RCHO; d) H<sub>2</sub>, Pd/C, MeOH; e) air oxidation.

**Figure 9** illustrates a method to synthesize compounds of Group III of this invention. The following reagents are used: a) LiBH<sub>4</sub>, THF; air oxidation; b) KH, ICH<sub>2</sub>CH<sub>2</sub>CH (OEt)<sub>2</sub>; then HCl, wet THF; c) thiazolidine-2,4-dione, piperidine, THF; d) H<sub>2</sub>, Pd/C, MeOH; e) air oxidation.

**Figure 10** illustrates a method to synthesize compounds of Formula II of the present invention.

**Figure 11** illustrates a method to synthesize compounds of Formula II of the present invention.

**Figure 12** illustrates a method to synthesize compounds of Formula II the present invention.

**Figure 13** illustrates a method to synthesize compounds of the present invention. The following reagents were used: a) LiBH<sub>4</sub>, THF; air oxidation; b) NaH, THF or DMF; c) thiazolidine-2,4-dione, piperidine, THF, or piperidine, benzoic acid; d) H<sub>2</sub>, Pd/C, MeOH; e) air oxidation.

**Figure 14** illustrates compounds of the present invention

**Figure 15** illustrates a method to synthesize compounds of the present invention: a) R or S or R/S. a) Lipoic Acid, DCC, Et<sub>3</sub>N; b) Method A: Zn, RCOX (*e.g.* acetic anhydride, succinic anhydride); Method B: NaBH<sub>4</sub>; then RCOX (*e.g.* XOCCH<sub>2</sub>NHt-BOC); c) When R is t-BOCNHCH<sub>2</sub> (*e.g.* tert-BOCglycine); deprotect with dry HCl in dioxane.

## GLOSSARY

As used herein, the term "alkyl" denotes branched or unbranched hydrocarbon chains, preferably having about 1 to about 8 carbons, such as, methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *sec*-butyl, *iso*-butyl, *tert*-butyl, octa-decyl and 2-methylpentyl. These groups can be optionally substituted with one or more functional groups which are attached commonly to such chains, such as, hydroxyl, bromo, fluoro, chloro, iodo, mercapto or thio, cyano, alkylthio, heterocyclyl, aryl, heteroaryl, carboxyl, carbalkoyl, alkyl, alkenyl, nitro, amino, alkoxyl, amido, and the like to form alkyl groups such as trifluoro methyl, 3--hydroxyhexyl, 2-carboxypropyl, 2-fluoroethyl, carboxymethyl, cyanobutyl and the like.

The term "alkylene" refers to a divalent alkyl group as defined above, such as methylene (-CH<sub>2</sub>-), propylene (-CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-), chloroethylene (-CHClCH<sub>2</sub>-), 2-thiobutene -CH<sub>2</sub>CH(SH)CH<sub>2</sub>CH<sub>2</sub>, 1-bromo-3-hydroxyl-4-methylpentene (-CHBrCH<sub>2</sub>CH(OH)CH(CH<sub>3</sub>)CH<sub>2</sub>-), and the like.

The term "alkenyl" denotes branched or unbranched hydrocarbon chains containing one or more carbon-carbon double bonds.

The term "alkynyl" refers to branched or unbranched hydrocarbon chains containing one or more carbon-carbon triple bonds.

The term "aryl" denotes a chain of carbon atoms which form at least one aromatic ring having preferably between about 6-14 carbon atoms, such as phenyl, naphthyl, and the like, and which may be substituted with one or more functional groups which are attached commonly to such chains, such as hydroxyl, bromo, fluoro, chloro, iodo, mercapto or thio, cyano, cyanoamido, alkylthio, heterocycle, aryl, heteroaryl, carboxyl, carbalkoyl, alkyl, alkenyl, nitro, amino, alkoxyl, amido, and the like to form aryl groups such as biphenyl, iodobiphenyl, methoxybiphenyl, anthryl, bromophenyl, iodophenyl, chlorophenyl, hydroxyphenyl, methoxyphenyl, formylphenyl, acetylphenyl, trifluoromethylthiophenyl, trifluoromethoxyphenyl, alkylthiophenyl,



trialkylammoniumphenyl, amidophenyl, thiazolylphenyl, oxazolylphenyl, imidazolylphenyl, imidazolylmethylphenyl, and the like.

The term "acyl" denotes the -C(O)R group, wherein R is alkyl or aryl as defined above, such as formyl, acetyl, propionyl, or butyryl.

5 The term "alkoxy" denotes -OR-, wherein R is alkyl.

The term "amido" denotes an amide linkage: -C(O)NR- (wherein R is hydrogen or alkyl).

The term "amino" denotes an amine linkage: -NR-, wherein R is hydrogen or alkyl.

10 The term "carboxyl" denotes -C(O)O-, and the term "carbonyl" denotes -C(O)-.

The term "carbonate" indicates -OC(O)O-.

The term "carbamate" denotes -NHC(O)O-, and the term "urea" denotes -NHC(O)NH-.

15 The term "EC<sub>50</sub>" refers to the concentration of a compound required to activate 50% of the receptors that bind the compound present in a sample or a subject. Thus, in the present invention, the EC<sub>50</sub> of a PPAR $\gamma$  modifier is the concentration of the modifier that activates 50% of the PPAR $\gamma$  present in the sample or organism. The term "activate" has its ordinary meaning, *i.e.*, cause to function or act.

20 The term "1,2-dithiolane" refers to a 5-membered heterocyclic ring consisting of two sulfur atoms at the 1 and 2 positions and carbon atoms at the remaining positions.

The term "dithiol" refers to a 1,3-dithiolpropanyl moiety.

25 The term "peroxisome proliferator activating receptor-gamma" or "PPAR $\gamma$ " refers to either the  $\gamma_1$ ,  $\gamma_2$  or  $\gamma_3$  isotypes or a combination of all isotypes of PPAR $\gamma$ . PPARs are nuclear receptors which naturally bind to fatty acids and which have been implicated in adipocyte differentiation (*see*, Perlmann & Evans, *Cell*, 90:391-397 (1997)).

30 The term "unit dosage form" refers to physically discrete units suitable as unitary dosages for human subjects and animals, each unit containing a predetermined quantity of active material calculated to produce the desired pharmaceutical effect in association with the required pharmaceutical diluent, carrier or vehicle. The specifications for the unit dosage forms of this invention are dictated by and dependent on

(a) the unique characteristics of the active material and the particular effect to be achieved and (b) the limitations inherent in the art of compounding such an active material for use in humans and animals. Examples of unit dosage forms are tablets, capsules, pills, powder packets, wafers, suppositories, granules, cachets, teaspoonfuls, tablespoonfuls, dropperfuls, ampoules, vials, aerosols with metered discharges, segregated multiples of any of the foregoing, and other forms as herein described.

The term "amino acid derivative" refers to an amino acid wherein the hydrogen on the  $\alpha$ -carboxylic acid function has been removed to generate a carboxyl group. Amino acids include, but are not limited to, the term amino acid as used herein refers to naturally occurring amino acids, amino acid analogs, and amino acid mimetics that function in a manner similar to the naturally occurring and analog amino acids. Naturally occurring amino acids are those encoded by the genetic code, as well as those amino acids that are later modified, *e.g.*, hydroxyproline,  $\gamma$ -carboxyglutamate, and O-phosphoserine. Amino acid analogs refers to synthetic amino acids that have the same basic chemical structure as a naturally occurring amino acid, *i.e.*, an  $\alpha$ -carbon that is bound to a hydrogen, a carboxyl group, an amino group, and an R group (*e.g.*, homoserine, norleucine, methionine sulfoxide, methionine methyl sulfonium). Such analogs have modified R groups (*e.g.*, norleucine) or modified peptide backbones, but retain the same basic chemical structure as a naturally occurring amino acid. Both naturally occurring and analog amino acids can be made synthetically by methods well known to those skilled in the art. Amino acid mimetics refer to chemical compounds that have a structure that is different from the general chemical structure of an amino acid, but that function in a manner similar to a naturally occurring amino acid.

The terms "cancer, neoplasm or malignancy" include primary and metastatic disease. So, for example, cervical cancer includes the neoplasm at the primary site (cervix) and metastatic cervical cancer, regardless of site of metastasis, such as skeleton, brain, *etc.*

The term "inflammatory disease" includes diseases (treatable or preventable with compounds described in this invention) including, but not limited to,

- a. T-lymphocyte activation and other T-lymphocyte-related disorders
- b. inflammatory cytokine (*e.g.* TNF- $\alpha$ , interleukin (IL)-1- $\alpha$ , IL-1- $\beta$ , IL-2, IL-6) production

c. activation of nuclear factors that promote transcription of genes encoding inflammatory cytokines. Examples of these nuclear transcription factors include but are not restricted to: nuclear factor-kappaB (NF-kappaB), activated protein-1 (AP-1), nuclear factor of activated T cells (NFAT)


5           The term "diabetes," unless stated or qualified otherwise, refers to all variant forms of diabetes mellitus (DM), including type 1 DM, type 2 DM, gestational diabetes, juvenile diabetes, *etc.*

          As used herein, the term "oxidative stress" refers to diseases or conditions that involve generation of active oxygen species and free radicals, resulting in the  
10 imposition of oxidative stress concomitant with the disease state. Examples of diseases imposing oxidative stress are dyslipidemias, diabetes mellitus and insulin resistant states, chronic viral infections (*e.g.* HIV, CMV, HSV, HBV, HCV infections), neurodegenerative diseases (*e.g.* Alzheimer's disease, multiple sclerosis, Parkinson's disease), cardiovascular disease (*e.g.* atherosclerosis, atherogenesis, vascular restenosis,  
15 congestive heart failure), diseases or conditions involving hypoxemia and hypoxic stress (stroke, vascular occlusive disease, MI, atherosclerosis, retinitis, retinal vein occlusion, hypoxic retinopathy, macular degeneration).

## 20                           **DETAILED DESCRIPTION OF THE INVENTION AND PREFERRED EMBODIMENTS**

### I.       **COMPOUNDS AND SYNTHESIS**

          This invention provides new thiazolindinedione derivatives. These  
25 compounds are useful for the treatment of certain cardiovascular diseases, certain endocrine diseases, certain inflammatory diseases, certain neoplastic (malignant), non-malignant proliferative diseases, certain neur-psychiatric disorders, certain viral diseases, and diseases associated with these viral infections as discussed herein. As such, in one aspect, the present invention provides a compound of Formula I:



wherein R, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup>, X, Y, n, m, q and t have been defined above. The bond bisecting the functional group containing two sulfurs, indicates that the bond can be attached at any of the 3 carbon atoms between the two sulfur atoms. The 1,2 dithiolane ring can be substituted at the 3-position or the 4-position. In a preferred embodiment, R<sup>1</sup> and R<sup>2</sup> are amino acid derivatives and thus, the compounds of Formula I are soluble in aqueous solution.

In the first preferred embodiment, Group I, R<sup>1</sup> and R<sup>2</sup> are hydrogen, or alternatively, R<sup>1</sup> and R<sup>2</sup> together with the sulfurs to which they are bound join to form a 1,2-dithiolane ring; X is O, NR, C(O)O, OC(O)O and C(O)NR, wherein R is a functional group including, but not limited to, hydrogen, optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl, and optionally substituted aryl.

Y is O, S and NR<sup>6</sup>, wherein R<sup>6</sup> is a member selected from the group consisting of hydrogen and optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl; n is an integer from 2 to 14; m is an integer from 1 to 6; q is 1; and t is 1. More preferably, n is 4, and X is COO, CONH and CON(CH<sub>3</sub>).

With reference to Figure 1, the first example to be discussed in Group I, is n is 4 and m is 2, *i.e.*, compound 12 (*see*, Figure 1). This is a preferred compound because it is believed that it will furnish a naturally occurring biochemical, lipoic acid 11, upon *in vivo* enzymatic hydrolysis.

The synthesis of analogs 12 and 13 begins with commercially available 4-hydroxybenzaldehyde 1 and 1,2-dibromoethane. Bromoethylation to furnish O-(2-bromoethyl)benzaldehyde 2 will occur under basic conditions (*see*, Nate, H. *et al.*, *Chem. Pharm. Bull.* 1987; 35(6): 2394-2411).

As for other chain lengths, *e.g.*, when m is 1, 3, 4, and 5, the starting benzaldehydes 6, 8-10 have been reported in the literature (*see*, Figure 2A). Aldehyde 6 has been described above along with 7. The three-carbon side chain 8, was described and prepared much in the same manner as the other bromides, *i.e.*, alkylation of the dihaloalkane with the anion of p-hydroxybenzaldehyde (*see*, Schweizer, E. *et al.*, *J. Org. Chem.* 1969; 34(1): 207-212). The four carbon aldehyde can be prepared according to the

method disclosed by Ito (*see*, Ito S, Japanese Patent No. 01,117,867). The ortho-analog of 10 has also been prepared (*see*, Nate, H. *et al.*, *Chem. Pharm. Bull.* 1987; **35**(6): 2394-2411).

With the bromoalkyl-benzaldehydes such as 7, S<sub>N</sub>2 alkylation with excess aqueous methylamine will provide the amine 3. Storage of 3 as its amine salt prevents self condensation reactions.

Two major approaches to the 5-benzylic-1,3-thiazolidine-2,4-dione ring system have been reported, (*see*, Figure 1, Yoshioka, T. *et al.*, U.S. Patent No. 4,572,912, Hulin, B. *et al.*, *J. Med. Chem.* 1992; **35**(10): 1853-1864, and Wilson, T. *et al.*, *J. Med. Chem.* 1996; **39**(3): 665-668).

Researchers recognized that the approach outlined in Figure 1 was cumbersome and thus, prepared euglycemic agents containing the thiazolidinedione ring system by an exceptionally mild Aldol condensation of thiazolidinedione itself to a benzaldehyde 14 (*see*, Figure 2B). The resulting arylidene thiazolidinedione 15 can then be reduced catalytically over Pd on C with hydrogen, or by the action of sodium-mercury amalgam to afford the desired ring system present in 16 (*see*, Figure 2B).

With reference to Figure 1, condensation of benzaldehyde 3 with thiazolidinedione and excess base will provide the 5-arylidene-thiazolidine-2,4-dione 4 that is easily reduced to provide the amine 5. Coupling of this secondary amine with R(+)-a-lipoic acid 11 can be achieved using standard activation of the carboxylate of 11 with species such as dicyclohexylcarbodiimide, which will lead to production of the target compound 12.

Reduction of the dithiolane group of lipoates has been reported using sodium borohydride, thus treatment of 12 with this reagent will give the potent antioxidant target dithiol 13 (*see*, Suzuki, Y.J. *et al.*, *Free Radical Res. Commun.* 1993; **18**: 115-122).

With reference to Figure 3, the example therein is illustrative of an ester tether, *i.e.*, X is C(O)O. Compound 20 is an example of this ester tether. Unlike the previous example of using an amino species 5 for coupling to lipoic acid 11, the corresponding alcohol is coupled. Hydroxyethylation of 1 can be accomplished under basic conditions using either ethylene oxide, or more conveniently, epichlorohydrin, as the electrophilic partner. Such reactions are common in phenol chemistry, for example, rutin has been exposed to epichlorohydrin in the presence of sodium hydroxide to give

hydroxyethylated rutin **23** (see, Figure 2C). (See, He, H. *et al.*, *Yiyao Gongye* 1987; **18**(5): 205-206). Preparation of the hydroxyethoxy derivative of quercetin is completed upon hydrolysis of the sugar moiety of **23** leading to **24**, wherein R = CH<sub>2</sub>CH<sub>2</sub>OH (see, Figure 2C).

5                   With reference to Figure 3, after the synthesis of **17**, Aldol condensation to give the thiazolidinedione **18** will be straightforward. Reduction of **18** in a manner outlined before is not complicated and will provide the alcohol **19**, ready for coupling to lipoic acid **11**. Coupling of the alcohol **19** with **11** will be a slower reaction than the amine **5** with **11** (Figure 1), and should a rate enhancement be desired,  
10   4-N,N'-dimethylaminopyridine (DMAP) will be used as a catalyst. Once the coupled target **20** is available, its reduction product **21** will be available by borohydride reduction of **20**.

                  With reference to Figure 4, it can be advantageous to employ an ether linkage in order to attach the lipoyl moiety to an aryl thiazolidinedione. Such an  
15   example, wherein X is oxygen, is exemplified by compound **30** wherein the natural number of carbon atoms present in lipoic acid are maintained in the molecule, and the chirality of the material is assured by starting with commercially available lipoic acid **11**. In order to make this attachment, it is necessary to reduce lipoic acid to the dithiolanyl alcohol **26**, and then attach this group by a Williamson ether synthesis to a benzaldehyde  
20   such as **7** as shown in Figure 4.

                  The most convenient manner of reducing lipoic acid is to reduce the methyl ester **25** with a strong reducing agent such as lithium borohydride, which will also reduce the dithiolane to a dithiol. However, during workup, air oxidation will lead to oxidation to the dithiolane **26**. With **26**, and the synthesis of **7** outlined above, coupling  
25   of **26** and **7** will occur under anhydrous conditions with NaH to give **27**. Aldol condensation to the thiazolidinedione **28** followed by reduction will provide the target dithiol **29**. Exposure of the target to ambient conditions will lead to facile air oxidation to target **30**.

                  In another preferred embodiment of Group I, X is a carbonate *i.e.*,  
30   OC(O)O. With reference to Figure 5, a carbonate linkage is illustrated in compound **34**, in which the antioxidant portion of the molecule is a reduced lipoate group. By starting with alcohol **26**, sequential addition of alcohols **26** and **17** to phosgene will provide the mixed carbonate **31**. Mild base catalyzed condensation of thiazolidinedione to the

aldehyde **31** will give **32**, reduction of which will give the target dithiol **33**. Air oxidation of **33** then provides the target **34**. Both targets can ultimately undergo *in vivo* metabolism of the carbonate group to furnish **26**. Ensuing enzymatic oxidations will readily transform **26** into lipoic acid.

5                    In another aspect, the present invention relates to compounds of Formula I wherein R<sup>1</sup> and R<sup>2</sup> are hydrogen, or alternatively R<sup>1</sup> and R<sup>2</sup> together with the sulfurs to which they are bound join to form a 1,2-dithiolane ring; X is O, NR, C(O)O, OC(O)O and C(O)NR, wherein R is a functional group including, but not limited to, hydrogen and optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl. Y is O, S and NR<sub>6</sub>, wherein R<sub>6</sub> is a member selected  
10 from the group consisting of hydrogen and optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl; n is an integer from 2 to 14; m is 0; q is 0; and t is 1. These are Group II compounds, and preferably, n is 4, and X is CONH or CON(CH<sub>3</sub>).

                    With reference to Figure 6, by excluding the tether adjacent to the aromatic ring, the target compounds appear shorter. Direct attachment via an amide bond  
15 of a benzaldehyde to lipoic acid is easily accomplished, and requires only a starting amino-benzaldehyde be obtained for coupling to lipoic acid. Specifically, **35** is available by numerous synthetic routes (*see*, Calmes, M. *et al.*, *Tetrahedron* 1990, **46**(17), 6021-6032 and Blokhin, A. V. *et al.*, *Khim. Geterotsikl. Soedin* 1990, **9**, 1226-1229).

                    Coupling of **35** to lipoic acid, as described earlier will provide **36**. If the  
20 active ester of lipoic acid and DCC are slow to react due to the lessened reactivity of the aniline, DMAP can be added to facilitate the reaction. After the synthesis of **36**, the subsequent sequence will follow from the earlier schemes. Aldol condensation to furnish the derivative **37** can then be followed by careful reduction and aerobic workup to provide **38**, which can be exposed to air to give the other desired target, **39**.

25                    In another preferred embodiment of Group II, X is oxygen. With reference to Figure 7, direct attachment of a reduced lipoic acid, *e.g.*, alcohol **26**, to a phenol (*e.g.*, **1**) will provide a stable class of thiazolidinedione-dithiolanes and only requires a reordering of events using intermediates already described. Thus, tosylation of **26** and rapid treatment with the phenoxide anion generated from 4-hydroxybenzaldehyde **1** will  
30 give aldehyde **40**. As before, once the correct benzaldehyde has been synthesized, elaboration of the thiazolidinedione moiety is straightforward. Aldol reaction of **40** to give **41**; reduction to **42** and finally, and finally air oxidation to generate **43**.

In yet another aspect, the present invention relates to compounds of Formula I wherein  $R^1$  and  $R^2$  are hydrogen, or alternatively  $R^1$  and  $R^2$  together with the sulfurs to which they are bound join to form a 1,2-dithiolane ring; X is O, NR, C(O)O, OC(O)O and C(O)NR, wherein R is a functional Group Including, but not limited to, hydrogen and optionally substituted ( $C_1$ - $C_6$ )alkyl. Y is O, S and NR<sub>6</sub>, wherein R<sub>6</sub> is a member selected from the group consisting of hydrogen and optionally substituted ( $C_1$ - $C_6$ )alkyl; n is an integer from 2 to 14; m is an integer from 2 to 14; q is 0; and t is 0. These are Group III compounds and preferably, n is 4 and X is C(O)NH, and C(O)N(CH<sub>3</sub>) as in lipoamides.

With reference to Figure 8, lipoic acid coupling with commercially available N-methyl-3-amino-propanol will generate the tether wherein X is C(O)N(CH<sub>3</sub>). Subsequent reduction of the alcohol with sodium hydride will yield the aldehyde. Mild base catalyzed condensation of thiazolidinedione to the aldehyde will give the intermediate product and reduction of which will give the target dithiol (49). Air oxidation then provides the target 1,2-dithiolane (50).

With reference to Figure 9, another preferred embodiment, is when X is oxygen. The most convenient manner of reducing lipoic acid is to reduce the methyl ester with a strong reducing agent such as lithium borohydride, which will also reduce the dithiolane to a dithiol. However, during workup, air oxidation will lead to oxidation to the dithiolane. Coupling of the dithiolane and ICH<sub>2</sub>CH<sub>2</sub>CH(OEt)<sub>2</sub> will occur under anhydrous conditions with potassium hydride. Aldol condensation with the thiazolidinedione followed by reduction will provide the target dithiol (59). Exposure of the target to ambient conditions will lead to facile air oxidation to target 1,2-dithiolane (60).

As will be apparent to those of skill in the art, some of the compounds of Formula I and II exist in optical, tautomeric, stereoisomeric and isomeric forms. For instance, when the 1,2 dithiothane is substituted at the 3-position, carbon number 3 can be in the R or S configuration. The compounds of the present invention encompass both the R and the S and any mixture in any percentage or racemic mixture thereof.


One aspect of the present invention relates to the use of various optical isomers of the lipoic acid derivatives. For example, in the case of  $\alpha$ -lipoic acid (R- and S-form, *i.e.* R-lipoic acid and S-lipoic acid), unlike the racemate (*see*, Biewenga *et al.* "An overview of lipoate chemistry." In: Lipoic Acid in Health and Disease., Fuchs J, Packer



L, Zimmer G, eds.), Marcel Dekker, Inc. 1997, pp 1-32, the R-enantiomer in certain instances has an anti-inflammatory activity, for example, being stronger by a factor of 10 than that of the racemate (Ulrich *et al.*, United States Patent 5,728,735 March 17, 1998). Moreover, in certain instances, the R-enantiomer has been shown to have superior  
5 insulin-sensitizing activity and to confer improved cardiac function (Zimmer G *et al.* *J.Mol Cel Cardiol.* 27:1895-903 (1995)) and capable of ameliorating diabetic peripheral and autonomic neuropathy (Ziegler D, Gries FA. *Diabetes.* 46 Suppl 2:S62-6 (1997)). In contrast, the S-enantiomer has been shown to be more effective as an antinociceptive agent. The antinociceptive (analgesic) activity of the S-enantiomer is for example  
10 stronger by a factor of 5 to 6 than that of the racemate (Ulrich *et al.*, United States Patent 5,728,735 March 17, 1998). Accordingly, the R- and S-enantiomers of the present invention are considered to have superior efficacy in the treatment of specific diseases. For example, the R-lipoic stereoisomeric thiazolidinedione derivatives have superior anti-inflammatory activity whereas the S-enantiomers provide greater efficacy in the treatment  
15 of other diseases.


Although these variations have been represented herein by a single molecular formula, the present invention includes the use of individual, isolated isomers, mixtures in various proportions and racemates thereof. Preferred compounds of Formula I are set forth in Table I below. Compounds of Formula I wherein R<sup>1</sup> and /or R<sup>2</sup> are  
20 amino acid derivatives, represent a preferred embodiment. Without being bound by any particular theory, it is believed that when Formula I has amino acid derivatives, the resulting derivatives are soluble in aqueous solution, and therefore, can act as prodrugs in biological systems.

TABLE I



COMP	R <sup>1</sup>	R <sup>2</sup>	n	X	m	Y	q	t
12	1,2 dithiolane		4	C(O)NCH <sub>3</sub>	2	O	1	1
13	H	H	4	C(O)NCH <sub>3</sub>	2	O	1	1
20	1,2 dithiolane		4	C(O)O	2	O	1	1
21	H	H	4	C(O)O	2	O	1	1
29	H	H	5	O	2	O	1	1
30	1,2 dithiolane		5	O	2	O	1	1
33	H	H	4	OC(O)O	2	O	1	1
34	1,2 dithiolane		4	OC(O)O	2	O	1	1
38	H	H	4	C(O)NCH <sub>3</sub>	0	-	0	1
39	1,2 dithiolane		4	C(O)NCH <sub>3</sub>	0	-	0	1
42	H	H	5	O	0	-	0	1
43	1,2 dithiolane		5	O	0	-	0	1
49	H	H	4	C(O)NCH <sub>3</sub>	3	-	0	0
50	1,2 dithiolane		4	C(O)NCH <sub>3</sub>	3	-	0	0
59	H	H	5	O	3	-	0	0
60	1,2 dithiolane		5	O	3	-	0	0
63	H	H	5	O	3	O	1	1
64	1,2 dithiolane		5	O	3	O	1	1
84	1,2 dithiolane		5	N-methyl	3	O	1	1
85	1,2 dithiolane		5	N-pyridyl	3	O	1	1
86	1,2 dithiolane		4	C(O)N	2	O	1	1
87	1,2 dithiolane		4	C(O)NCH <sub>3</sub>	0	-	0	-
88	1,2 dithiolane		1	C(O)NCH <sub>3</sub>	2	O	1	1
89	1,2 dithiolane		5	C(O)NCH <sub>3</sub>	2	O	1	1
90	1,2 dithiolane		5	C(O)N salt	2	O	1	1

In another aspect, the present invention relates to a compound of Formula II:



5 wherein R, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup>, X, Y, m, n and q have been defined above. The bond bisecting the functional group containing two sulfurs indicates that the bond can be attached at any of the 3 carbon atoms between the two sulfur atoms. In an alternative embodiment, R<sup>1</sup> and R<sup>2</sup> together with the sulfurs to which they are bound join to form a 1,2-dithiolane ring. The 1,2 dithiolane ring can be substituted at the 3-  
10 position or at the 4-position.

In this aspect of the invention, compounds of Formula II can be prepared in similar fashion as outlined above for compounds of Formula I. In Formula II, structures retain the lipoate residues, but a carboxylic acid or ester replaces the thiazolidinedione ring. Compounds of Formula II have similar PPAR-γ activity and  
15 similar biological effects as do compounds of Formula I, but have altered pharmacokinetics and metabolism. The acids *i.e.*, wherein R<sup>5</sup> is a hydrogen, could be administered as ester prodrugs (*i.e.*, wherein R<sup>5</sup> is alkyl), thus enhancing their bioavailability.

With reference to Figure 10, beginning with arylacetic acids and their  
20 esters such as 65 wherein X' is a halogen, alkylation of dibromoalkanes of varying carbon chain-length will lead to formation of bromoalkylethers 66. Upon formation of the anion of alcohol 26 and exposure to the bromides 66, alkylated products like 67 will be generated. The reaction will work for the dianion (wherein R<sup>5</sup> is hydrogen in 65) thus requiring 2 equivalents of base to deprotonate both phenol and carboxylate. In either  
25 case, the phenoxide anion is the more reactive anion towards the dihalide, leading to 66. If 67 is produced as the ester, it can be hydrolyzed, or if 67 is produced as the acid, it can be esterified. The dithiol 68 can be produced from 67 by simple reduction in the absence of air.


With reference to Figure 11, a sulfur atom can replace the phenolic linking  
30 atom. Conversion of the phenol 65 to a thiophenol 69 is possible using dimethylthiocarbonylchloride (*see*, Newman, *et al.*, *Organic Synthesis*, Vol. 51, pg 139,

1971). The ensuing chemistry follows with minor modification to provide thioethers **71** or **72** wherein  $R^5$  is hydrogen or alkyl; and where  $X'$  is hydrogen or halogen.

An alternative procedure of linkage of lipoic acid to either **65** or **69** to give **74** or **75** (*see*, Figure 12) is possible. Either thioester **74** or ester **75** is metabolized *in vivo* to naturally occurring lipoic acid with regeneration of the acids/esters **65** and/or **69**. The synthesis of either **74** or **75** requires simple esterification of **65** or **69** with the active ester of lipoic acid **73** ( $X'$  is Cl, OCOOEt, *etc.*) in the presence of a base such as pyridine or a trialkylamine. These active esters are formed directly from lipoic acid by any number of standard methods such as DCC, pyridine; oxalyl chloride, NaH; ethyl chloroformate and triethylamine, or the like.

In certain other aspects, compounds of Formula II also include other attachment methods such as esters, amides, or thioester linkages. In the case wherein  $X$  in Formula II is a ester, the previous intermediate halides **70** or **66** can be reacted with the carboxylate anion of **73** ( $X'$  is O-Metal) to give products **76** or **77** directly. In certain aspects, wherein  $X$  in Formula II is an amide, the halides are first converted to an amine or thiol by simple  $S_N2$  chemistry. Thus, reaction of **70**, for example, with  $LiNHR'$  gives the amine **78** that can then be reacted with **73** to furnish **76**. Alternatively, **79** can be reacted with **73** to give **76** wherein  $X$  in Formula II is a thioester.

In yet another aspect, the present invention relates to a compound of Formula III:



In Formula III,  $R^1$  and  $R^2$  are each independently a functional group including, but not limited to, hydrogen,  $C(O)-R^6$  and  $C(S)-R^6$ .  $R^6$ , in Formula III, is a functional group including, but not limited to, hydrogen,  $(C_1-C_{12})$ alkyl, aryl, arylalkyl,  $(C_1-C_{12})$ carboxyl,  $(C_1-C_{12})NHR^7$ ,  $(C_1-C_{12})NR^7R^8$ ,  $OR^7$ ,  $NHR^7$ ,  $SR^7$ ,  $NR^7R^8$ .  $R^7$  and  $R^8$ , in Formula III, are each independently a functional group including, but not limited to, hydrogen,  $(C_1-C_{12})$ alkyl, aryl and arylalkyl. The bond bisecting the functional group containing two sulfurs indicates that the bond can be attached at any of the 3 carbon atoms between the two sulfur atoms.

In an alternative embodiment, R<sup>1</sup> and R<sup>2</sup> together with the sulfurs to which they are bound join to form a 1,2-dithiolane ring. The 1,2 dithiolane ring can be substituted at the 3-position or at the 4-position.

X, in Formula III, is functional group including, but not limited to O, NR, C(O)O, OC(O)O and C(O)NR, wherein R is a functional group including, but not limited to, hydrogen and optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl. Y, in Formula III, is a functional group including, but not limited to, O, S and NR<sup>6</sup>, wherein R<sup>6</sup> is a functional group including, but not limited to, hydrogen and optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl. In Formula III, the index "n" is an integer from 2 to 14; the index "m" is an integer from 0 to 14; the index "q" is an integer from 0 to 1; the index "r" is an integer from 0 to 4; and the index "t" is an integer from 0 to 1, provided when m is 0 then q is 0.

## II. CHARACTERIZATION AND PURIFICATION OF THE TARGETS

The synthetic chemistry outlined above can be carried out by standard methods apparent to those skilled in the art, and employ purification of reaction products by chromatography and/or crystallization. Product homogeneity can be ascertained by high performance liquid chromatography. A variety of columns (normal phase silica gel, reverse phase C-18, etc.) are available, as are computer workstations to analyze the results. Once reaction products are deemed greater than 99.5% HPLC pure, they can be analyzed by elemental analysis, NMR spectroscopy, FTIR, UV and EI or CI mass spectroscopy. Exact mass determinations will be possible and particularly applicable to intermediates. Other physical properties can be determined and recorded such as solubility, melting point, stability, *etc.* A careful study of chemical stability can be performed and suitable formulation for the oral route of administration can be examined.

## III. BIOLOGICAL ASSAY

Compounds of Formulae I, II and III are activators of PPAR $\gamma$ . As described hereinbelow, a transient cotransfection assay can be used to screen for PPAR $\gamma$  antagonist. In this assay, chimeras are constructed that fuse the ligand binding domains of three murine PPAR subtypes ( $\alpha$ ,  $\gamma$  and NUC-1) to the DNA binding domain of the yeast transcription factor GAL4 (*see* Example 18). Expression plasmids for the GAL4-PPAR chimeras are then transfected into CV-1 cells with a reported construct containing five copies of the GAL4 DNA binding site upstream of the thymidine kinase (tk) promoter

driving chloramphenicol acetyl transferase (CAT) gene expression. Using this assay system, compounds of Formulae I and II which are activators of PPAR $\gamma$  and not PPAR $\alpha$  and NUC-1 are identified. (see, J.M. Lehmann *et al.*, *J. Biol. Chem.* 270:12953-12956 (1995)).


5                   Plasmids - GAL4-PPAR chimera expression constructs contain the translation initiation sequence and amino acids 1-76 of the glucocorticoid receptor fused to amino acids 1-147 of the yeast transcription factor GAL4, including the DNA binding domain, in the pSG5 expression vector (Stratagene). cDNAs encoding amino acids 167-468, 138-440, and 174-475 of murine PPAR $\alpha$ , NUC-1, and PPAR $\gamma$ 1 are amplified by  
10 polymerase chain reaction and inserted C-terminal to GAL4 in the pSG5 expression vector (Stratagene) to generate plasmids pSG5-GAL4-PPAR $\alpha$ , pSG5-GAL4-NUC-1, and pSG5-GAL4-PPAR $\gamma$ , respectively. The regions of the PPARs included in the chimeras should contain the ligand binding domains based on their homology to ligand binding domains of characterized nuclear receptors. The chimeras initially contained the  
15 translation start site and N-terminal 262 amino acids of the glucocorticoid receptor, including the  $\tau$ 1 transcriptional transactivation domain. However, as these chimeras had high basal activity in CV-1 cells, a 0.6-kilobase BglII fragment containing the  $\tau$ 1 domain can be removed, leaving the translation start site and amino acids 1-76 of the glucocorticoid receptor. Wild-type receptor expression vectors can be generated by  
20 insertion of cDNAs encoding murine PPAR $\alpha$ , NUC-1, PPAR $\gamma$ 1 and PPAR $\gamma$ 2 into the expression vector pSG5 (Stratagene). Reporter plasmid (UAS)5-tk-CAT can be generated by insertion of five copies of a GAL4 DNA binding element into the BamHI site of pBLCAT2. The reporter a P2-tk-CAT was generated by insertion of the 518-bp EcoRI/XBAI fragment containing the enhancer of the aP2 gene into the BamHI site of  
25 pBLCAT2.

*Cotransfection Assay* -- CV-1 cells are plated in 24-well plates in DME medium supplemented with 10% delipidated fetal calf serum. In general, transfection mixes contain 10 ng of receptor expression vector, 100 ng of the reporter plasmid, 200 ng of  $\beta$ -galactosidase expression vector (pCH110, Pharmacia) as internal control, and 200 ng  
30 of carrier plasmid. Transfections can be done with Lipofectamine (Life Technologies, Inc.) according to the manufacturer's instructions. Cell extracts were prepared and assayed for chloramphenicol acetyltransferase and  $\beta$ -galactosidase activities as described previously.

*Ligand Binding Assay* -- cDNA encoding amino acids 174-475 of PPAR $\gamma$ 1 can be amplified via polymerase chain reaction and inserted into bacterial expression vector pGEX-2T (Pharmacia). GST-PPAR $\gamma$  LBD can be expressed in BL21(DE3)plysS cells and extracts prepared as described previously. For saturation binding analysis,  
 5 bacterial extracts (100  $\mu$ g of protein) is incubated at 4°C for 3 h in buffer containing 10 mM Tris (pH 8.0), 50 mM KCl, 10 mM dithiothreitol with [ $^3$ H]-BRL49653 (specific activity, 40 Ci/mmol) in the presence or absence of unlabeled BRL49653. Bound can be separated from free radio-activity by elution through 1-ml Sephadex G-25 desalting  
 10 columns (Boehringer Mannheim). Bound radioactivity eluted in the column void volume and can be quantitated by liquid scintillation counting.


#### IV. COMPOSITIONS AND METHODS

In another aspect, the present invention provides a pharmaceutical composition comprising a compound of Formula I:



wherein R, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup>, X, Y, n, m, q and t have been defined above, or a pharmaceutically acceptable salt or solvate thereof, and a pharmaceutically acceptable carrier.

20 In yet another aspect, the present invention provides a pharmaceutical composition comprising a compound of Formula II:



25 wherein R, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup>, X, Y, m, n and q have been defined above, or a pharmaceutically acceptable salt or solvate thereof, and a pharmaceutically acceptable carrier. Suitable salts include, but are not limited to, sodium, potassium, ammonium, and calcium.

The compounds of this invention can be formulated in a variety of carriers and delivery systems. For instance, to prepare a long-acting depot formulation, a therapeutically effective concentration of the compound is placed in an oil, resin,

biopolymer or other suitable delivery device as is known in the art. The amount of the therapeutic compound to be administered and the compound's concentration in depot formulations depend upon the vehicle or device selected, the clinical condition of the patient, the side effects and the stability of the compound in the formulation. Thus, the physician employs the appropriate preparation containing the appropriate concentration of the therapeutic compound and selects the amount of formulation administered, depending upon clinical experience with the patient in question or with similar patients.

In addition to the therapeutic compound, the compositions can include, depending on the formulation desired, pharmaceutically-acceptable non-toxic carriers, or diluents which include vehicles commonly used to form pharmaceutical compositions for animal or human administration. The diluent is selected so as not to unduly affect the biological activity of the combination. Examples of such diluents which are especially useful for injectable formulations are water, the various saline solutions, Ringer's solution, dextrose solution, and Hank's solution. In addition, the pharmaceutical composition or formulation may include additives such as other carriers; adjuvants; or nontoxic, non-therapeutic, non-immunogenic stabilizers and the like.

Furthermore, excipients can be included in the formulation. Examples include cosolvents, surfactants, oils, humectants, emollients, preservatives, stabilizers and antioxidants. Any pharmacologically acceptable buffer may be used, e.g., Tris or phosphate buffers. Effective amounts of diluents, additives and excipients are those amounts which are effective to obtain a pharmaceutically acceptable formulation in terms of solubility, biological activity, *etc.*

Thus, a composition of the invention includes a therapeutic compound which can be formulated with conventional, pharmaceutically acceptable, vehicles for topical, oral or parenteral administration. Formulations can also include small amounts of adjuvants such as buffers and preservatives to maintain isotonicity, physiological and pH stability. Means of preparation, formulation and administration are known to those of skill. *See generally Remington's Pharmaceutical Science*, 15th ed., Mack Publishing Co., Easton, Pa. (1980).

To prepare a topical formulation for the treatment of dermatological disorders listed in Tables II, III, IV & V, or diseases of the external eye (Table VII), a therapeutically effective concentration of the compound is placed in a dermatological vehicle as is known in the art. The amount of the therapeutic compound to be




administered and the compound's concentration in the topical formulations depend upon the vehicle selected, the clinical condition of the patient, the side effects and the stability of the compound in the formulation. Thus, the physician employs the appropriate preparation containing the appropriate concentration of the therapeutic compound and  
 5 selects the amount of formulation administered, depending upon clinical experience with the patient in question or with similar patients.

The concentration of the therapeutic compound for topical formulations is in the range of about 1 mg/mL to about 1000 mg/mL. Typically, the concentration of the therapeutic compound for topical formulations is in the range of about 2.5 mg/mL to  
 10 about 25 mg/mL. Solid dispersions of the therapeutic compound as well as solubilized preparations can be used. Thus, the precise concentration is subject to modest experimental manipulation in order to optimize the therapeutic response. About 2,500 mg of therapeutic compound per 100 grams of vehicle is useful in the treatment of skin lesions to provide a 2.5% weight/weight (w/w) formulation. Suitable vehicles include  
 15 oil-in-water or water-in-oil emulsions using mineral oils, petrolatum and the like as well as gels such as hydrogel.


Alternative topical formulations include shampoo preparations, oral paste, and mouth wash preparations. ORABASE7 can be used as the base oral paste to which the therapeutic compound is added. Concentrations of therapeutic compound are  
 20 typically as stated above for topical formulations.

In yet another aspect, the present invention relates to a method of treating a PPAR $\gamma$  mediated disease or oxidative stress, comprising administering a therapeutically effective amount of compound of the Formula I:



25 wherein R, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup>, X, Y, n, m, q and t have been defined above, to an individual suffering from a PPAR $\gamma$  mediated disease.

In still yet another aspect, the present invention relates to a method of treating a PPAR $\gamma$  mediated disease, comprising administering a therapeutically effective amount of compound of the Formula II



wherein R, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup>, X, Y, m, n and q have been defined above, to an individual suffering from a PPAR $\gamma$  mediated disease.

Diseases treatable with the novel compounds described herein are outlined in Tables II, III, IV, V, VI & VII. These compounds have clinical utility in the treatment of non-malignant (Table II) and malignant (Table III) diseases in multiple organ systems, diseases caused by naked or coated DNA and RNA viruses, and local and disseminated diseases associated with the infection by these viruses (Table IV), human immunodeficiency virus (HIV) infection and diseases associated with HIV infection (Table V), neuro-psychiatric diseases (Table VI), and diseases of the eye (Table VII).

The methods of treatment provided by this invention are practiced by administering to a human or vertebrate animal in need a dose of a 1,2-dithiolane derivative compound that binds to or modifies the activity of peroxisome proliferator activated receptor-gamma (PPAR $\gamma$ ), or a pharmaceutically acceptable salt or solvate thereof. The present method includes both medical therapeutic and/or prophylactic treatment as necessary.

The compounds described in this invention can be use to treat a variety of disorders including proliferative, inflammatory, metabolic, or infectious disorders. The specific disorders that can be treated with the compounds described in this invention are listed in Tables II, III, IV, V, VI & VII.

Using methods of the invention, therapeutic compounds are typically administered to human patients topically or orally. Parenteral administration is used in appropriate circumstances apparent to the practitioner. Preferably, the compositions are administered in unit dosage forms suitable for single administration of precise dosage amounts. For example long-acting depot compositions are administered subcutaneously or intra-muscularly as precise unit doses with each dose lasting weeks to months.

## V. ADMINISTRATION

The therapeutic compound is optionally administered topically by the use of a transdermal therapeutic system (*see*, Barry, *Dermatological Formulations*, (1983) p. 181 and literature cited therein). While such topical delivery systems have been designed

largely for transdermal administration of low molecular weight drugs, by definition they are capable of percutaneous delivery. They can be readily adapted to administration of the therapeutic compounds of the invention by appropriate selection of the rate-controlling microporous membrane.

5                   For ophthalmic applications (Table VII), the therapeutic compound is formulated into solutions, suspensions, and ointments appropriate for use in the eye. The concentrations are usually as discussed above for topico-local preparations. For ophthalmic formulations, see Mitra (ed.), *Ophthalmic Drug Delivery Systems*, Marcel Dekker, Inc., New York, N.Y. (1993) and also Havener, W. H., *Ocular Pharmacology*,  
10                   C.V. Mosby Co., St. Louis (1983).

                  The therapeutic compound is alternatively administered by aerosol. This is accomplished by preparing an aqueous aerosol, liposomal preparation or solid particles containing the compound. A nonaqueous (e.g., fluorocarbon propellant) suspension could be used. Sonic nebulizers are preferred because they minimize exposing the therapeutic  
15                   compound to shear, which can result in degradation of the compound.

                  Ordinarily, an aqueous aerosol is made by formulating an aqueous solution or suspension of the therapeutic compound together with conventional pharmaceutically acceptable carriers and stabilizers. The carriers and stabilizers vary with the requirements of the particular compound, but typically include nonionic surfactants (Tweens,  
20                   Pluronic, or polyethylene glycol), innocuous proteins like serum albumin, sorbitan esters, oleic acid, lecithin, amino acids such as glycine, buffers, salts, sugars or sugar alcohols. Aerosols generally are prepared from isotonic solutions.

                  For oral administration, either solid or fluid unit dosage forms can be prepared. For preparing solid compositions such as tablets, the compound of interest is  
25                   mixed into formulations with conventional ingredients such as talc, magnesium stearate, dicalcium phosphate, magnesium aluminum silicate, calcium sulfate, starch, lactose, acacia, methylcellulose, and functionally similar materials as pharmaceutical diluents or carriers. Capsules are prepared by mixing the compound of interest with an inert pharmaceutical diluent and filling the mixture into a hard gelatin capsule of appropriate  
30                   size. Soft gelatin capsules are prepared by machine encapsulation of a slurry of the compound of interest with an acceptable vegetable oil, light liquid petrolatum or other inert oil. Fluid unit dosage forms for oral administration such as syrups, elixirs and suspensions can be prepared. The water soluble forms can be dissolved in an aqueous

vehicle together with sugar, aromatic flavoring agents and preservatives to form a syrup. An elixir is prepared by using a hydroalcoholic (*e.g.*, ethanol) vehicle with suitable sweeteners such as sugar and saccharin, together with an aromatic flavoring agent. Suspensions can be prepared with an aqueous vehicle with the aid of a suspending agent  
5 such as acacia, tragacanth, methylcellulose and the like.

Appropriate formulations for parenteral use are apparent to the practitioner of ordinary skill. Usually, the therapeutic compound is prepared in an aqueous solution (discussed below) in a concentration of from about 1 to about 100 mg/mL. More typically, the concentration is from about 10 to 60 mg/mL or about 20 mg/mL.

10 Concentrations below 1 mg/mL may be necessary in some cases depending on the solubility and potency of the compound selected for use. The formulation, which is sterile, is suitable for various parenteral routes including intra-dermal, intra-articular, intra-muscular, intravascular, and subcutaneous.

Slow or extended-release delivery systems, including any of a number of  
15 biopolymers (biological-based systems), systems employing liposomes, and polymeric delivery systems, can be utilized with the compositions described herein to provide a continuous or long term source of therapeutic compound. Such slow release systems are applicable to formulations for topical, ophthalmic, oral, and parenteral use.

## 20 VI. ROUTES OF ADMINISTRATION

Therapeutic agents of the invention can be delivered or administered topically or by transdermal patches for treating disorders involving the skin. Oral administration is preferred for disorders in Tables II, III, IV, V, VI & VII that cannot be treated effectively by topical therapy. Additionally, the agents can be delivered  
25 parenterally for the conditions in listed in Tables II, III, IV, V, VI & VII, that do not respond to oral or topical therapy or for conditions where oral or topical therapy is not feasible. Parenteral therapy is typically intra-dermal, intra-articular, intramuscular or intravenous.

A preferred way to practice the invention for disorders in II, III, IV, V, &  
30 VII that affect the skin is to apply the compound of interest, in a cream, lotion, ointment, or oil based carrier, directly to the skin lesions. Typically, the concentration of therapeutic compound in a cream, lotion, or oil is 1-2%. Alternatively, an aerosol can be used topically. These compounds can also be orally administered.

In general, the preferred route of administration is oral, parenteral, or topical (including administration to the eye, scalp, and mucous membranes). Topical administration is preferred in treatment of skin lesions, including lesions of the scalp, lesions of the cornea (keratitis), and lesions of mucous membranes where such direct application is practical. Shampoo formulations are sometimes advantageous for treating scalp lesions such as seborrheic dermatitis and psoriasis of the scalp. Mouthwash and oral paste formulations can be advantageous for mucous membrane lesions, such as oral lesions and leukoplakia.

Oral administration is a preferred alternative for treatment of dermatological and eye diseases listed in Tables II, III, IV, V, VI & VII where direct topical application is not useful, and it is a preferred route for other non-dermatological applications. Intravenous administration may be necessary in disorders that cannot be effectively treated by topical or oral administration. Intra-articular injection is a preferred alternative in cases of arthritis (psoriatic or nonpsoriatic) where the practitioner wishes to treat one or only a few (such as 2-6) joints. Usually, the compound is delivered in an aqueous solution of about 10-20 mg/mL. Additionally, the therapeutic compounds are injected directly into lesions (intra-lesion administration) in appropriate cases. Intradermal administration is an alternative for dermal lesions.

For pulmonary applications, a chemical delivery system for drug targeting to lung tissue using a compound of Formulae I, II or III or mixtures thereof as the "targetor moiety". Therefore a preferred therapeutic compound is compound 3 that modifies the activity of PPAR $\gamma$  and is formulated into solutions, suspensions, aerosols and particulate dispersions appropriate for application to the pulmonary system. The therapeutic agent may be inhaled via nebulizer, inhalation capsules, inhalation aerosol, nasal solution, intratracheal as a solution via syringe, or endotracheal tube as an aerosol or via as a nebulizer solution. In vitro kinetic and in vivo pharmacokinetics studies have shown that the compounds of Formulae I, II or III or mixtures thereof, provide an effective pulmonary delivery system which, in a sufficiently stable in buffer and biological media, is hydrolyzed rapidly into the respective active parent drugs, with significantly enhanced delivery and retention of the active compound to lung tissue.

## VII. DOSAGE AND SCHEDULES

An effective quantity of the compound of interest is employed in treatment. The dosage of compounds used in accordance with the invention varies depending on the compound and the condition being treated. The age, weight, and clinical condition of the recipient patient; and the experience and judgment of the clinician or practitioner administering the therapy are among the factors affecting the selected dosage. Other factors include the route of administration the patient, the patient's medical history, the severity of the disease process, and the potency of the particular compound. The dose should be sufficient to ameliorate symptoms or signs of the disease treated without producing unacceptable toxicity to the patient.

Broadly, an oral dosing schedule is from about 100 to about 600 mg twice a day. More typically, a single dose is about 100-200 mg of compound given twice a day. A convenient oral dose for an adult patient is 200 mg twice a day. A dosage range for topical treatment is about 0.1% to about 10% (weight/volume) in a cream or ointment, applied twice a day. A usual dose for intra-articular injection is 20-40 mg injected per joint, not generally exceeding three joints per therapy session. A typical dosage for intra-dermal administration is about 20-75 mg per injection per site. A typical dosage for intravenous or intramuscular administration in an adult patient would be between 1 mg and 1000 mg per day given in single or divided doses depending on the judgment of the practitioner.

In some aspects the oral dose is determined from the following formula:

$$\text{Oral dose (in milligrams)} = (k_1)(EC_{50})(k_2)(LBW)(MW);$$

wherein  $k_1$  is a dimensionless constant with values ranging from 5 to 100;  $EC_{50}$  is in mol/L;  $k_2$  is the fractional water content of the lean body weight (LBW) of the patient = 0.72 L/kg, (see, *GEIGY SCIENTIFIC TABLES*, VOL 1, Lentner (ed.), p217, Ciba-Geigy Limited, Basle, Switzerland (1981); and MW is the molecular weight of the drug in g/mol.

Typically, the dosage is administered at least once a day until a therapeutic result is achieved. Preferably, the dosage is administered twice a day, but more or less frequent dosing can be recommended by the clinician. Once a therapeutic result is achieved, the drug can be tapered or discontinued. Occasionally, side effects warrant discontinuation of therapy. In general, an effective amount of the compound is that which

provides either subjective relief of symptoms or an objectively identifiable improvement as noted by the clinician or other qualified observer.

The compounds in this invention can also be given orally in combination with compounds that bind or modify the activity of the vitamin D receptor or in  
5 combination with compounds that bind or modify the activity of the retinoid X receptors or retinoic acid receptors to provide for a synergistic effect in the treatment or prevention of the disorders listed in Tables II, III, IV, V, VI & VII. Examples of such compounds that provide for synergistic effect when given in combination with the drugs encompassed  
10 by the current invention include vitamin D analogs, various retinoic acid derivatives, and other ligands for retinoid X receptors or retinoic acid receptors including but not limited to compounds such as LG100268, tazarotene, TTNPB, or LGD1069 (Targretin), and RXR/PPAR heterodimeric ligands, *e.g.* LG101754 (Ligand Pharmaceuticals) (see, United States Patent No. 6,004,987 Demarchez, *et al.*, December 21, 1999, Use of ligands which are specific for RXR receptors.).

15 In certain aspects, the compounds in the invention can also be given systemically (oral administration preferred) or topically in combination with natural or synthetic ligands (agonist or antagonist ligands) that bind and modify the activity of the estrogen (nuclear) receptor (ER), which include the alpha and/or the beta isoforms, *i.e.* ER-alpha or ER-beta or both ER-alpha and ER-beta. Examples of ER-alpha and/or ER-  
20 beta ligands are: estradiol, selective (synthetic) estrogen receptor modulators such as raloxifene, anti-estrogens such as tamoxifen and hydroxytamoxifen, and the naturally occurring phytoestrogens such as the isoflavones genistein, diadzein and glycitein.

In other aspects, the compounds of the present invention are given independently for uses specified herein, or are given as adjuvants, *i.e.* in combination  
25 with existing pharmacological therapeutic agent or agents or other pharmaceutical agent or agents yet to be discovered. For example, when used as anti-cancer therapy for the treatment of breast cancer, the compounds in this invention are given either independently or as adjuvant therapy in combination with one or more other anti-cancer agent(s) efficacious in the treatment of breast cancer, *e.g.* tamoxifen, megestrol acetate,  
30 cyclophosphamide, methotrexate, 5-fluorouracil, doxorubicin, prednisone). Another example is the use of the compounds in this invention for the prevention of allograft rejection, either independently or as adjuvant therapy in combination with one or more other immunosuppressive agent(s). Examples of these include immunosuppressive

glucocorticoids (*e.g.* prednisone), inhibitors of purine synthesis (*e.g.* azathioprine and mycophenolate), and inhibitors of the calcineurin-dependent cytokine synthesis in activated lymphocytes (*e.g.* cyclosporine, tacrolimus and sirolimus). (see, United States Patent No. 5,925,657, Seed, *et al.* July 20, 1999, Use of PPAR. $\gamma$  agonists for inhibition of inflammatory cytokine production.).

Synergistic therapeutic effects can be achieved by oral administration of the drugs encompassed in the current invention together with orally or intravenously administered drugs that bind to and or modify the activity of either the vitamin D receptor or retinoid X receptors or retinoic acid receptors. As such, in another embodiment, the present invention relates to a method of treating a PPAR $\gamma$  mediated disease, comprising administering a combination therapy of a compound of Formulae I, II or III and a member selected from the group consisting of a drug that bind to or modifies the activity of a vitamin D receptor, a retinoid X receptor, or a retinoic acid receptor.

A preferred dosage range for administration of a retinoic acid derivative or retinoid would typically be from 0.1 to 100 mg per square-meter of body surface area, depending on the drug's ability to bind to or modify the activity of its cognate nuclear receptor, given in single or divided doses, orally or by continuous infusion, two or three times per day. For synergistic therapy, the preferred dosages and routes and frequency of administration of the vitamin D analogs or retinoid compounds can be similar to the dosages and routes and frequency of administration ordinarily recommended for these agents when given without compounds of Formulae I, II or III. Examples of effective retinoids are 9-cis-retinoic acid, 13-cis-retinoic acid, all-trans-retinoic acid (at-RA). Preferred retinoids for this purpose would include 13-cis-retinoic acid, tazarotene, or Targretin. A preferred dosage range for systemic administration of a vitamin D analog would typically be from 0.1 to 100 mg per square-meter of body surface area, depending on the drug's ability to bind to and or activate its cognate vitamin D receptor, given in single or divided doses, orally or by continuous infusion, two or three times per day. Examples of effective vitamin D analogs are 1,25-dihydroxy-vitamin D (1,25-(OH) $_2$ -vit D) and calcipotriene. The dosage range and routes and frequency of administration of compounds of Formulae I, II or III required to achieve synergistic effects when given with vitamin D or retinoid derivatives are the same as those described elsewhere in this disclosure. The preferred mode of administration of these drugs for synergistic therapeutic purposes would be orally although alternatively one can use topical or



parenteral routes of administration. Synergistic therapeutic effects can also be achieved for conditions that are treated by topical administration of vitamin D derivatives or retinoid related compounds such as psoriasis, acne, or other disorders not involving the skin described in Tables II, III, IV, V, VI & VII. The dosages and the modes and  
5 frequency of administration of the vitamin D or retinoid related compounds for synergistic topical therapy would be similar to those ordinarily recommended for these agents when given without compounds of Formulae I, II or III. The dosage range and the modes and frequency required for topical administration of the compounds of Formulae I, II or III given in combination with vitamin D or retinoid related compounds are the same  
10 as those described elsewhere in this disclosure.

The foregoing is offered primarily for purposes of illustration. It will be readily apparent to those of ordinary skill in the art that the operating conditions, materials, procedural steps and other parameters of the system described herein may be further modified or substituted in various ways without departing from the spirit and  
15 scope of the invention. For example, the invention has been described with human patients as the usual recipient, but veterinary use is also contemplated.

## VIII. EXAMPLES

### General

20 All reactions were carried out under an argon atmosphere with dry, freshly distilled solvents under anhydrous conditions, unless otherwise stated. Acetone was distilled from potassium carbonate, and N, N-dimethylformamide (DMF), from calcium hydride. Yields were applied to chromatographically and spectroscopically (<sup>1</sup>H-NMR) homogeneous materials. Reagents were purchased at highest commercial quality and  
25 used without further purification unless otherwise mentioned. Reactions were monitored by thin-layer chromatography carried out on 250 microns Analtech, Inc. silica gel plates, employing UV light and p-anisaldehyde solution and heat to visualize the distributions of compounds. Silica gel (60 Å, 230-400 mesh) (Whatman Inc.) was used for flash column chromatography. Preparative thin-layer chromatography (PTLC) separations were  
30 carried out on 1000 microns silica gel plates (Analtech). NMR spectra were obtained by Bruker Advance DXR-400 or DXR-300 instruments and calibrated using tetramethylsilane. Abbreviations were used to explain multiplicities as followings: s = singlet, d = doublet, t = triplet, q = quartet, qui = quintet, m = multiplet, b = broad.

### Example 1

This example illustrates the synthesis of compounds **63** and **64**

A. The synthesis of compound **8**.

5 With reference to Figure 13,  $K_2CO_3$  (1.82 g, 13 mmol) and 1,3-dibromopropane (3.05 mL, 30 mmol) were added to a solution of 4-hydroxy-benzaldehyde (1.22 g, 10 mmol) in acetone (30 mL). The mixture was heated under reflux for 8.5 hrs. Then the reaction mixture was filtered and the solvent was removed by evaporation. The resulting residue was mixed with ether and then washed consecutively  
10 with 5% NaOH, water, and brine. After drying with  $MgSO_4$  and evaporation of the solvent, flash column chromatography (silica gel, hexane 100% > EtOAc : hexane = 10 : 90 > 20 : 80) gave the desired compound as a pale yellow oil:  $R_f$  = 0.71 (silica gel, EtOAc : hexane = 35 : 65); yield 61%;  $^1H$ -NMR (300 MHz,  $CDCl_3$ )  $\delta$  9.86 (1H, s), 7.82-7.79 (2H, dd,  $J$  = 6.9 and 1.8 Hz), 7.00-6.97 (2H, dd,  $J$  = 7.1 and 1.6 Hz), 4.19-4.15 (2H, t,  $J$  = 5.8 Hz), 3.61-3.56 (2H, t, 6.4 Hz), 2.37-2.28 (2H, qui, 6 Hz).

B. The synthesis of compound **61**.

With reference to Figure 13, to a methyl lipoate (**25**) solution in dry THF (10 grams or 45 mmol, in 500 mL) at 0 °C under argon was added a 1M  $LiBH_4$  solution in  
20 THF (50 mmol or 50 mL). After the reduction was complete by TLC, the mixture was acidified with dilute HCl in water, and extracted 3 X 150 mL diethyl ether. The combined organic layer was dried over  $NaSO_4$ , and the solvent was removed by rotary evaporation to provide **26** as a thick oil. Upon treatment of a suspension of NaH (1.4 grams of a 48% oil dispersion, washed free of oil with hexane, 30 mmol) in dry THF (50  
25 mL) under argon with **26** in dry THF (5 grams or 26 mmol in 100 mL) hydrogen gas is evolved with formation of the sodium salt of **26**. Addition of a THF solution of the bromobenzaldehyde **8** prepared in part A above, (30 mmol in 50 mL) leads to formation of sodium bromide and the desired product. Water is added (1 L) and the mixture is extracted 3 X 100 mL ethylacetate. The combined organic layer was dried over  $NaSO_4$ ,  
30 filtered, and the solvent evaporated to give the aldehyde **61**. The aldehyde is then purified by crystallization or column chromatography.

C. Synthesis of the Aldol adduct compound **62**.


A solution of the benzaldehyde **61** (5 grams or 14 mmol) in ethanol (100 mL) was treated with piperidine (0.5 mL) and heated at reflux. Alternatively, the benzaldehyde could be treated in benzene with piperidinium benzoate (0.5 grams) with azeotropic removal of water. In either case, when the reaction was judged complete by  
5 TLC, the reaction mixture is poured into water and extracted with EtOAc (3X100 mL); the combined organic layer is then dried over NaSO<sub>4</sub>, filtered, and the solvent removed to give the Aldol adduct **62**.

D. Synthesis of the 1,2 dithiolane compound **64**.

10 The purified Aldol adduct **62** (2 grams or 4.4 mmol) in ethanol (50 mL) is treated with excess 10% Pd/C (5 g), or with a similar quantity of S-resistant Pd/C, and placed under an atmosphere of hydrogen with stirring. When reduction was complete by TLC, the Pd/C was removed by filtration under argon, and the solvent was evaporated  
15 generate the dithiolane **64** that could be purified by crystallization or chromatography.

**Example 2**

This example illustrates the synthesis of compound **86** N-(2-{4-[(2,4-dioxo(1,3-thiazolidin-5-yl)methyl]phenoxy}ethyl)-5-(1,2-dithiolan-3-yl) pentanamide.




20 In a 50 mL round bottomed flask was taken 0.102g (0.3 mmol) of hydrochloride salt of 5-({4-[2-aminoethoxy]phenyl}methyl)-1,3-thiazolidine-2,4-dione in 10 mL of dry dichloromethane. 0.05 mL (0.3 mmol) of triethylamine was added at 0°C and the mixture was stirred at this temperature for an additional 10 minutes. 0.068g  
25 (0.3 mmol) of dl- lipoic acid was dissolved in 10 mL of dry dichloromethane in a second round bottomed flask which was cooled to 0°C and to this was added 0.05 mL (0.3 mmol) of triethylamine followed by 0.321 ml (0.3 mmol) of isopropyl chloroformate (1M solution in toluene) dropwise over 10 mins. The mixture was stirred for an additional 10 min at this temperature and cannula transferred to the first flask containing the free  
30 amine. After 1 hr 50 ml water was added and the aqueous layer was extracted with 2\*

50 mL of dichloromethane. The combined organic layer was washed with 10% sodium bicarbonate and brine. The organic phase was dried over sodium sulfate and concentrated under vacuum. The residue was chromatographed over silica (eluent: 35% ethyl acetate: hexane) to yield 0.064g (56%)  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ): 8.3 (bs, 1H); 7.14 (d,  $J = 8.4\text{Hz}$ , 2H); 6.83 (d,  $J = 8.4\text{Hz}$ , 2H); 5.9 (bs, 1H); 4.5 (dd,  $J = 4\text{Hz}$ , 9.3Hz, 1H); 4.02 (t,  $J = 5\text{Hz}$ , 2H); 3.6 (dd, 2H); 3.5 (m, 1H); 3.4 (dd,  $J = 4\text{Hz}$ , 16Hz, 1H); 3.2- 3 (m, 3H); 2.3 (m, 2H); 2.2 (t,  $J = 7.4\text{Hz}$ , 1H); 1.8 (m, 2H); 1.75-1.5 (m, 6H), 1.5-1.45 (m, 2H)  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ): 25.7, 29.2, 30.3, 34.9, 36.7, 38, 38.8, 39.3, 40.6, 54, 56.7, 67.2, 115.1, 128, 130.9, 157, 170, 173, 174. HRMS: (Cal for  $\text{C}_{20}\text{H}_{26}\text{N}_2\text{O}_4\text{S}_3$ ) Calculated 455.1127, Found 455.1096 IR ( $\text{cm}^{-1}$ ): 3021, 2926, 1752, 1698, 1609, 1516, 1303, 1244, 1154, 1052.

### Example 3

This example illustrates the synthesis of N-(2-{4-[(2,4-dioxo(1,3-thiazolidin-5-yl)methyl]phenoxy}ethyl)-5-(1,2-dithiolan-3-yl)-N-methylpentanamide compound 12.



In a 250 mL round bottomed flask was taken 0.802g (2.6 mmol) of hydrochloride salt of 5-({4-[2-(methylamino)ethoxy]phenyl}methyl)-1,3-thiazolidine-2,4-dione in 30 mL of dry dichloromethane and to it was added 0.370 mL (2.6 mmol) of triethylamine at  $0^\circ\text{C}$  stirred at this temperature for an additional 10 minutes. 0.542g (2.7 mmol) of dl- lipoic acid was dissolved in 50 mL of dry dichloromethane in a second round bottomed flask which was cooled to  $0^\circ\text{C}$  to this was added 0.370 mL (2.6 mmol) of triethylamine followed by 2.6 mL (2.6 mmol) of isopropyl chloroformate (1M solution in toluene) dropwise over 10 mins. After 10 minutes this mixture was cannula transferred to the first flask containing the free amine and stirred at  $0^\circ\text{C}$  for 1 hr. Water was added at the end of 1 hr and the aqueous layer was extracted with 2\* 50 mL of dichloromethane. The combined organic layer was washed with 10% sodium bicarbonate and brine. The dichloromethane was removed under vacuum and residue chromatographed over silica (eluent MeOH:  $\text{CHCl}_3$ , 1: 100) to yield 0.891g (72%)  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ): 8.19 (bs, 1H); 7.15 (d,  $J = 8\text{Hz}$ , 2H); 7.13 (d,  $J = 8.6\text{Hz}$ , 2H); 6.83 (d,  $J = 8.6\text{Hz}$ , 2H); 6.82 (d,  $J = 8\text{Hz}$ ,


2H); 4.49 (dd,  $J = 5\text{Hz}$ ,  $12.3\text{Hz}$ , 1H); 4.1 (t,  $J = 6.7\text{Hz}$ , 2H); 3.47 (t,  $J = 6.7\text{Hz}$ , 2H); 3.56 (m, 1H); 3.42 (dd,  $J = 5\text{Hz}$ ,  $18\text{Hz}$ , 1H); 3.1 (s, 1H); 3.1 (m, 2H); 2.44 (m, 2H); 2.3 (t,  $J = 7.4\text{Hz}$ , 1H); 1.89 (m, 2H); 1.65 (m, 4H), 1.45 (m, 2H).

$^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ): 24.3(t), 24.7(t), 28.7(t), 28.8(t), 32.4 (t), 32.9(t), 33.9(q), 34.6(t),  
 5 37.5(q), 37.6(t), 38.3(d), 39.9(d), 47.7(t), 48.9(t), 53.3(d), 56.3(d), 65(t), 66.3(t), 114.5(d),  
 128.1(s), 128.5(s), 130.2(d), 130.4 (d), 157.5(s), 157.9(s), 170.7(s), 173.3(s), 173.4(s),  
 174.5(s). HRMS: (Cal for  $\text{C}_{21}\text{H}_{28}\text{N}_2\text{O}_4\text{S}_3$ ) Calculated 469.1284, Found 469.1263 IR  
 (film,  $\text{cm}^{-1}$ ): 3030, 2926, 1752, 1696, 1609, 1512, 1303, 1244, 1154, 1052.

10

### Example 4


This example illustrates the preparation of N-{4-[(2,4-dioxo(1,3-thiazolidin-5-yl)methyl]phenyl}-5-(1,2-dithiolan-3-yl)-N-methyl pentanamide compound  
 87.



15 To a solution of 0.072g (0.3 mmol) of amine in 2 mL of dry dichloromethane was added 0.062g (0.32 mmol) of dl-lipoic acid followed by 0.062g (0.3 mmol) of dicyclohexylcarbodiimide at  $0^\circ\text{C}$ . The resulting mixture was stirred at room temperature for 3 hrs. The precipitated dicyclohexylurea was filtered and washed with 5 mL of dichloromethane. The filtrate was concentrated under vacuum and the residue was  
 20 chromatographed over silica gel (eluent ethyl acetate : hexanes, 3: 7) to yield 0.098g (76%).  $^1\text{H}$ NMR ( $\text{CDCl}_3$ ): 8.7 (bs, 1H); 7.29 (d,  $J = 8.15\text{Hz}$ , 2H); 7.15 (d,  $J = 8.15\text{Hz}$ ); 4.54 (dd,  $J = 4\text{Hz}$ ,  $9\text{Hz}$ , 1H); 3.53 (dd,  $J = 4\text{Hz}$ ,  $14\text{Hz}$ , 1H); 3.49 (m, 1H); 3.25 (s, 3H); 3.22-3.07 (m, 3H); 2.41 (dddd,  $J = 6.2\text{Hz}$ ,  $18.7\text{Hz}$ , 1H); 2.04 (t, 2H); 2.05 (dddd,  $J = 6.2\text{Hz}$ ,  $18.6\text{Hz}$ , 1H); 1.58 (m, 4H); 1.32 (m, 2H).  $^{13}\text{C}$ NMR ( $\text{CDCl}_3$ ): 25.5(t), 29.2(t),  
 25 34.2(t), 34.9(t), 37.2(d), 38.5(t), 38.8(t), 40.6(t), 53.3(d), 56.8(q), 128.0(d), 131.1(d), 136.1(s), 143.8(s), 170.6(s), 173.5(s), 174.6(s). HRMS (Calculated for  $\text{C}_{19}\text{H}_{24}\text{N}_2\text{O}_3\text{S}_3$ ): Calculated 425.1022, Found 425.1007 IR (film,  $\text{cm}^{-1}$ ): 2925, 2852, 1753, 1699, 1624, 1601, 1511, 1461, 1390.

**Example 5**

This example illustrates the preparation of N-(2-{4-[(2,4-dioxo(1,3-thiazolidin-5-yl)methyl]phenoxy}ethyl)-5-(1,2-dithiolan-3-yl)-N-methylacetamide compound **88**.



5


In a 50 mL round bottomed flask was taken 0.144g (0.472 mmol) of hydrochloride salt of 5-({4-[2-(methylamino)ethoxy]phenyl}methyl)-1,3-thiazolidine-2,4-dione in 10 mL of dry dichloromethane. 0.065 mL (0.472 mmol) of triethylamine was added at 0°C and the mixture was stirred at this temperature for an additional 10 minutes. 0.077g (0.472 mmol) of 2-(1,2-dithiolan-3-yl)acetic acid was dissolved in 10 mL of dry dichloromethane in a separate round bottomed flask which was again cooled to 0°C. To this was added 0.065 mL (0.472 mmol) of triethylamine followed by 0.469 mL (0.472 mmol) of isopropyl chloroformate (1M solution in toluene) dropwise over 10 mins. The mixture was stirred for an additional 10 min at this temperature. The mixed anhydride thus formed was canula transferred to the other flask containing the free amine. The combined mixture was stirred at 0°C for 1 hr. Water was added and the aqueous layer was extracted with 2\* 50 mL of dichloromethane. The combined organic layer was washed with 10% sodium bicarbonate and brine. The dichloromethane was removed under vacuum and residue chromatographed over silica (eluent: 30% ethyl acetate: hexane) to yield 0.084g (56%) <sup>1</sup>H NMR (CDCl<sub>3</sub>): 8.6 (bs, 1H); 7.14 (d, *J* = 8.4Hz, 2H); 6.83 (d, *J* = 8.42, 2H); 4.48 (dd, *J* = 4Hz, 9Hz, 1H); 4.12 (t, *J* = 5Hz, 1H); 3.75 (t, 2H), 3.6 (m, 2H); 3.1 (s, 3H); 3.0 (dd, *J* = 4Hz, 16Hz, 1H); 2.8-2.65 (m, 2H); 2.6 (m, 1H); 2.0 (m, 1H). <sup>13</sup>C NMR (CDCl<sub>3</sub>): 34.4, 37.9, 38.1, 39.7, 40, 40.3, 48.5, 50.8, 51.2, 54, 65.4, 66.8, 115, 128.6, 130.8, 130.9, 157.9, 158.3, 171.5, 171.7, 171.9, 175.2. HRMS (Calculated for C<sub>18</sub>H<sub>22</sub>N<sub>2</sub>O<sub>4</sub>S<sub>3</sub>) Calculated 427.0814, Found 427.0807 IR (film, cm<sup>-1</sup>): 3025, 2924, 2850, 1749, 1695, 1634, 1611, 1510, 1405, 1302, 1243, 1154.

20

25

## Example 6

This example illustrates the synthesis of preparation of 5-[(4-{2-[5-(1,2-dithiolan-3-yl)pentyl)methylamino]ethoxy}phenyl)methyl]-1,3-thiazolidine-2,4-dione compound 89.



5

To a solution of 4.8g (10mmol) of N-(2-{4-[(2,4-dioxo(1,3-thiazolidin-5-yl)methyl]phenoxy}ethyl)-5-(1,2-dithiolan-3-yl)-N-methylpentanamide in 100 mL of dry tetrahydrofuran at 0°C was added 40 mL of alane-N,N-dimethylethylamine complex (20 mmol) dropwise over 0.5hr. After stirring the reaction mixture at that temperature for an additional 1 hr, 20 mL of water was added and the mixture was stirred for 10 min. The precipitated aluminum hydroxide was filtered and washed with ethyl acetate. The filtrate was mixed and the tetrahydrofuran removed under vacuum. The residue was partitioned between ethyl acetate and water and the aqueous phase extracted 2\*50 mL of ethyl acetate. The combined ethyl acetate layer was washed with brine and concentrated to leave dithiol as an oil. The residue was dissolved in 40 mL of ethyl acetate and cooled to 0°C. 100 mL of chilled 10% sodium bicarbonate was added followed by 2.6g (10 mmol) of iodine dissolved in 50 mL ethyl acetate dropwise till brown color persists. After a few minutes 10% sodium thiosulphate solution was added to destroy excess iodine. The ethyl acetate layer was separated and the aqueous layer was extracted with 2\*75 mL of ethyl acetate. The combined ethyl acetate layer was washed with brine dried over sodium sulfate and concentrated to leave a residue which was chromatographed over silica gel (eluent MeOH: CHCl<sub>3</sub>, 2: 100) to yield 1.86g (47% based on recovered starting) along with 0.865g of starting amide.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) : 7.2 (d, J = 8.5Hz, 2H); 6.77 (d, J = 8.5Hz, 2H); 4.34 (dd, J = 3.7Hz, 9Hz, 1H); 4.06 (t, J = 5.4Hz, 2H); 3.5 (m, 1H); 3.4 (dd, J = 3.6Hz, 18Hz, 1H); 3.2-3.0 (m, 3H); 2.9 (t, J = 5.3Hz, 2H); 2.58 (dd, J = 9.0Hz, 18Hz, 1H); 2.44 (dddd, J = 6.7Hz, 18Hz, 1H); 2.4 (s, 3H), 1.89 (dddd, J = 6.7Hz, 18Hz, 1H); 1.6 (m, 2H); 1.5 (m, 2H), 1.45 (m, 2H); 1.34 (m, 2H).

<sup>13</sup>C NMR (CDCl<sub>3</sub>) : 25.4(t), 27.4(t), 29.4(t), 35.1(t), 38.4(t), 38.8 (t), 40.6(t), 41.7(d), 55(d), 55.5(t), 56.9(q), 57.4(t), 65(t), 114.9(d), 129.7(s), 130.8(d),


30

157.7(s), 175.5(s), 180(s). HRMS (Calculated for  $C_{21}H_{30}N_2O_3S_3$ ) Calculated 455.1491, Found 455.1507 IR (film,  $cm^{-1}$ ): 2925, 2851, 1749, 1697, 1607, 1586, 1511, 1462, 1301, 1243.

5

### Example 7

This example illustrates the preparation of tartrate salt of 5-[(4-{2-[5-(1,2-dithiolan-3-yl)pentyl)methylamino]ethoxy}phenyl)methyl]-1,3-thiazolidine-2,4-dione compound 90.



10


To 1.2g (2.6mmol) of 5-[(4-{2-[5-(1,2-dithiolan-3-yl)pentyl)methylamino]ethoxy}phenyl)methyl]-1,3-thiazolidine-2,4-dione dissolved in 50 mL of dry tetrahydrofuran at 0°C was added 0.396g (2.6 mmol) of dl-tartaric acid dissolved in 40 mL of dry tetrahydrofuran. After the addition was complete the mixture was stirred at that temperature for 12 hrs. The tetrahydrofuran was removed under vacuum and the residue was washed with chloroform to remove unreacted amine and recrystallized from ether: pentane to yield 1.48g (92%)  $^1H$  NMR ( $DMSO-d_6$ ) : 7.15 (d,  $J$  = 8.4, 2H); 6.88 (d,  $J$  = 8.4Hz, 2H); 4.82 (dd,  $J$  = 4.2Hz, 9Hz, 1H); 4.19 (t, 2H); 4.13 (s, 2H); 3.5 (m, 1H); 3.3 (dd,  $J$  = 4Hz, 16Hz, 1H); 3.16 (t, 2H); 3.1-2.9 (m, 3H); 2.8 (m, 2H); 2.57 (s, 3H); 2.38 (m, 1H); 1.83 (m, 1H), 1.7-1.4 (m, 4H); 1.2-1.4 (m, 4H).  $^{13}C$  NMR ( $DMSO-d_6$ ) : 25.1(t), 26(t), 29.2(t), 35(t), 37.2(t), 38.9(t), 40.7(t), 41.6(d), 54(d), 55(t), 56.7(t), 57(q), 64.3(t), 72.9(d), 115.2(d), 130(s), 131.2(d), 157.7(s), 172.8(s), 174.8(s), 174.9(s), 177(s).

25

### Example 8

This example illustrates the preparation of N-(2-{4-[(2,4-dioxo(1,3-thiazolidin-5-yl)methyl]phenoxy}ethyl)-N-methyl-6,8-disulfanyloctamide.






To a solution of 0.100g (0.21 mmol) N-(2-{4-[(2,4-dioxo(1,3-thiazolidin-5-yl)methyl]phenoxy}ethyl)-5-(1,2-dithiolan-3-yl)-N-methylpentanamide in 2 mL of tetrahydrofuran at 0°C was added 0.016g (0.4 mmol) of sodium borohydride dissolved in 0.5 mL water. The mixture was allowed to stir at that temperature for an additional 1 hr. 2 mL 10% HCl was then added and the tetrahydrofuran removed under vacuum. The residue was extracted with 2\*50 mL of chloroform. The combined chloroform layer was washed with brine, dried over sodium sulfate and then concentrated under vacuum. The residue was chromatographed over silica gel (eluent 1% methanol in chloroform) to yield 0.089mg (86%) <sup>1</sup>H NMR (CDCl<sub>3</sub>) : 10.4 (bs, 1H); 7.08 (d, *J* = 8Hz, 2H); 6.76 (d, *J* = 8Hz, 2H); 4.39 (dd, *J* = 9Hz, 4Hz, 1H); 4.09 (t, *J* = 5.6 Hz, 2H); 3.73 (t, *J* = 5.6Hz, 2H); 3.42 (dd, *J* = 4Hz, 16Hz, 1H); 3.14 (s, 3H); 3.12 (dd, *J* = 16Hz, 9Hz, 1H); 2.8 (m, 1H); 2.6 (m, 2H); 2.33 (t, *J* = 9.7Hz, 1H); 1.8 (m, 1H); 1.7- 1.4 (m, 6H); 1.3-1.36 (m, 2H); <sup>13</sup>C NMR (CDCl<sub>3</sub>) : 22.7(t), 24.9(t), 25.3(t), 27(t), 27.2(t), 33.2(t), 33.7(t), 34.5(d), 38(q), 38.19(t), 39.1(d), 40.6(q), 43.1(t), 48.4(t), 49.5(t), 56.8(t), 65.6(t), 66.9(t), 115(d), 128.8(s), 129.2(s), 130.7(d), 130.8(d), 157.9(s), 158.2(s), 171.6(s), 171.7(s), 174.1(s), 174.3(s), 175.5(s), 175.6(s); HRMS (calculated for C<sub>21</sub>H<sub>30</sub>N<sub>2</sub>O<sub>4</sub>S<sub>3</sub>) Calculated 471.1440, Found 471.1442; IR (film, cm<sup>-1</sup>): 3032, 2926, 1752, 1698, 1609, 1513, 1303, 1244, 1154, 1055.

20

### Example 9

This example illustrates the preparation of 6,8-diacetylthio-N-(2-{4-[(2,4-dioxo(1,3-thiazolidin-5-yl)methyl]phenoxy}ethyl)-N-methyloctanamide.



0.075g (0.164mmol) of N-(2-{4-[(2,4-dioxo(1,3-thiazolidin-5-yl)methyl]phenoxy}ethyl)-N-methyl-6,8-disulfanyloctanamide was dissolved in 1 mL pyridine and to it was added 0.034g (0.33mmol) of acetic anhydride. The mixture was stirred for 2 hrs at room temperature. 10 mL 10% HCl was added and the product extracted with 2\*50 mL of chloroform. The combined chloroform layer was washed with

brine dried over sodium sulfate and concentrated to leave crude diacetate which was further purified by column chromatography over silica gel (eluent 1% methanol: chloroform) to yield 0.88g;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ) : 8.9 (bs, 1H); 7.14 (d,  $J = 8.3\text{Hz}$ , 2H); 7.12 (d,  $J = 8.3\text{Hz}$ , 2H); 6.8 (d,  $J = 8.3\text{Hz}$ , 2H); 6.7 (d,  $J = 8.3\text{Hz}$ , 2H); 4.7 (dd,  $J = 4\text{Hz}$ , 9.3Hz); 4.09 (t,  $J = 5.2\text{Hz}$ , 2H); 3.72 (t,  $J = 5.2\text{Hz}$ , 2H); 3.6 (m, 1H); 3.4 (dd,  $J = 4\text{Hz}$ , 16Hz, 1H); 3.13 (s, 3H); 3.12 (dd,  $J = 9.3\text{Hz}$ , 16Hz, 1H); 3-2.7 (m, 2H); 2.3 (m, 6H); 2.3 (t,  $J = 6\text{Hz}$ , 2H); 1.9-1.7 (m, 2H); 1.7-1.5 (m, 4H); 1.5- 1.2 (m, 2H).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ) : 25, 25.2, 26.9, 27, 31, 31.2, 32.6, 33.7, 34.2, 35, 35.1, 37.9, 38.1, 44.0, 48.3, 49.2, 54, 54.1, 66, 67, 115, 128.5, 129, 130.7, 130.9, 157, 158.4, 171.2, 173.8, 174, 175, 196.1, 196.2. HRMS (calculated for  $\text{C}_{25}\text{H}_{34}\text{N}_2\text{O}_6\text{S}_3$ ) : Calculated 555.1652, Found 555.1746 IR (film,  $\text{cm}^{-1}$ ): 3032, 2936, 1752, 1742, 1698, 1609, 1513, 1309, 1244, 1154, 1052.

### Example 10

This example illustrates a clinical trial and therapy by topical application.

15 A patient having dermal manifestations of either psoriasis vulgaris, or acne vulgaris, or human papilloma virus (HPV) infection (*e.g.*, anogenital warts) is selected for therapy using the invention. A compound of Formulae I, II or III that modifies the activity of  $\text{PPAR}\gamma$  is prepared in a cream vehicle at a concentration of 1 to 5% (weight/volume), typically 2.5% and is applied to the affected skin three times a day.

20 After the skin lesions have subsided, therapy is discontinued.

### Example 11

This example illustrates a clinical trial and therapy by oral administration.

25 A patient having type 2 diabetes mellitus, or chronic generalized acne, or chronic generalized psoriasis, with or without psoriatic arthritis, or rheumatoid arthritis, or inflammatory bowel disease (*e.g.*, ulcerative colitis) is selected for therapy. The patient weighs 80 kilograms. For infants or children the doses suggested are lowered in a linear fashion based on body weight or surface area. The female patient of child-bearing potential is given a pregnancy test to confirm that the patient is not pregnant. Provided

30 that the patient is not pregnant and does not plan to become pregnant during treatment, a compound of Formulae I, II or III that modifies the activity of  $\text{PPAR}\gamma$  is orally administered in a dosage of 20 to 1,000 milligrams twice daily, more typically 100 mg twice daily. The patient is monitored for improvement in the manifestations of the index

disease. Additionally, a complete blood count, including white cell count and differential, a platelet count, and liver function tests (such as levels of alkaline phosphatase, lactose dehydrogenase, and transaminases) are checked prior to treatment and periodically thereafter. The dosage is tapered when the manifestations of the disease subside, or  
5 discontinued if indicated.

### Example 12

This example illustrates a clinical trial and therapy by intravenous injection administration.

10 A patient having non-metastatic cancer, such as breast cancer, prostate cancer or colon cancer is selected for therapy. The patient weighs 80 kilograms. For infants or children the doses suggested are lowered in a linear fashion based on body weight or surface area. The female patient of child-bearing potential is given a pregnancy test to confirm that the patient is not pregnant. If indicated, the tumor is surgically  
15 excised. When the patient is stable post-surgically, a compound of Formulae I, II or III that modifies the activity of PPAR $\gamma$  administered intravenously in a dose of 50 to 1,000 mg, more typically in a dosage of 200 milligrams, as a bolus or continuous infusion over 4 hr, every 12 hr. The patient is monitored for improvement in his or her manifestations of the cancer in terms of laboratory tests such as prostate specific antigen (PSA) for  
20 prostatic cancer, or carcinoembryonic antigen (CEA) for breast or colon cancer. Additionally, a complete blood count, including white cell count and differential, a platelet count, and liver function tests (such as levels of alkaline phosphatase, lactose dehydrogenase, and transaminases) are checked prior to treatment and periodically thereafter. The dosage is tapered when the manifestations of the disease subside, or  
25 discontinued if indicated.

### Example 13

This example illustrates a clinical trial and therapy by intralesional injection administration.

30 The patient is one who has venereal warts (HPV infection), or dermatological manifestations of Kaposi sarcoma (with or without infection with the human immunodeficiency virus) who is not a candidate for surgery, or in whom surgery may impair bodily functions (such as painful sexual activity, or impairment in urination

or defecation in the case of venereal warts). The patient is treated by administering, intralesionally, injection(s) of a compound of Formulae I, II or III that modifies the activity of PPAR $\gamma$ . The compound is administered in a 5 to 50 mg/mL, more typically 20 mg/mL of an aqueous solution, a suspension or an emulsion. About 10-75 mg are  
5 injected directly into each lesion, depending on its size and volume. The therapy is repeated weekly until the lesions are eradicated.

#### Example 14

This example illustrates a clinical trial and therapy by intra-articular  
10 injection administration.

A patient having psoriatic arthritis or rheumatoid arthritis with painful, swollen, inflamed joints, is treated with intra-articular injection(s) of a compound of Formulae I, II or III that modifies the activity of PPAR $\gamma$ . The compound is administered in a 5 to 50 mg/mL, more typically 20 mg/mL of an aqueous solution, a suspension or an  
15 emulsion. About 10-50 mg are injected directly into each joint, depending on the joint and the severity of the disease. The therapy is repeated weekly until the disease subsides and the pain and inflammation resolves.

#### Example 15

This example illustrates a clinical trial and therapy by intrathecal injection  
20 administration.

A patient having viral meningitis (*e.g.*, meningitis caused by cytomegalovirus), is treated with intrathecal injection(s) of a compound of Formulae I, II or III that modifies the activity of PPAR $\gamma$ . The compound is administered by injection or  
25 via catheter, in a 5 to 50 mg/mL (more typically 20 mg/mL) of an aqueous solution, a suspension or an emulsion. About 10-50 mg, depending on the patient's lean body-mass and the severity of the disease, is injected directly into the intrathecal space. The therapy is administered 3 time a week and tapered to once weekly as symptoms subside, and eventually discontinued when the infectious agent is eradicated.

30

#### Example 16

This example illustrates a clinical trial for prophylaxis or therapy of ocular disease by topical application.

For ophthalmic applications, the therapeutic compound is a compound of Formulae I, II or III that modifies the activity of PPAR $\gamma$  and is formulated into solutions, suspensions, and ointments appropriate for application to the external eye. The formulation contains the drug at a concentration of 20 mg/mL. A patient having allergic conjunctivitis, or viral conjunctivitis, or keratitis, corneal ulceration or uveitis (secondary to surgical procedures or contact lens injury) is treated by topico-local administration of the optic formulation, with 2 to 3 drops being instilled in each eye and the process repeated every 4 hr for 2 wk or more, depending on the type or severity of the disease, until the symptoms have subsided or resolved.

10

### Example 17

This example illustrates a clinical trial for pulmonary therapy.

A compound of Formulae I, II or III is used and is formulated into solutions, suspensions, aerosols and particulate dispersions appropriate for application to the pulmonary system. The patient weighs 80 kilogram. For infants or children the doses suggested are lowered in a linear fashion based on body weight or surface area. The patient has asthma or COPD. The therapeutic agent may be inhaled via nebulizer, inhalation capsules, inhalation aerosol, nasal solution, intratracheal as a solution via syringe, or endotracheal tube as an aerosol or via as a nebulizer solution. For delivery via nebulizer, the therapeutic agent is given at a dose of 0.5 to 50 mg per 2 mL solution, more typically 10 mg/2 mL dose given every 15 min as needed for acute treatment. For delivery by metered dose inhaler, the therapeutic agent is given at a dose of 0.05 to 1.0 mg per actuation, more typically 0.2 mg per actuation, given every 4 hr.

15

20

COMPOUND 12 is a potent activator of PPAR $\gamma$ .

25

### Example 18

This example illustrates activating PPAR $\gamma$  with compounds of the present invention.

30

African green monkey kidney cells were co-transfected with: 1) a PPAR $\gamma$  construct in which the DNA binding domain of yeast GAL4 sequence was fused to the ligand binding domain of PPAR $\gamma$  and 2) a reporter construct with 4 copies of the upstream activation sequence of GAL4 fused to a luciferase reporter gene. Cells were

incubated with varying concentrations of compound 12, rosiglitazone, or troglitazone and activation of the luciferase reporter construct was then recorded using a luminometer after addition of substrate. All three thiazolidinediones tested were capable of activating PPAR $\gamma$  with compound 12 being far more potent than even rosiglitazone.

5

### Example 19

This example illustrates compounds of the present invention are antidiabetic.

Male insulin resistant obese Zucker rats were treated by oral administration of compound 12 in a dose of 100 mg/kg/day for one week. The drug was suspended in a mixture of 0.5% carboxymethylcellulose and administered by gavage. Male insulin resistant obese Zucker rats treated with vehicle not containing compound 12 were used as placebo controls. After one week, blood samples were obtained for measurement of serum insulin levels by radioimmunoassay and serum glucose levels by glucose oxidase method. The percent reduction in serum insulin and glucose levels induced by administration of compound 12 was determined by comparing results in treated animals versus placebo controls. The effects of a higher dose of orally administered Rezulin® (troglitazone) on serum insulin and glucose levels in male obese Zucker rats as reported by Fujiwara *et al.*, *Diabetes* 37:1549 are shown for comparison purposes. Administration of compound 12, like administration of a higher dose of troglitazone, was associated with reductions in serum insulin levels and glucose levels. The reductions in glucose levels in the face of reductions in insulin levels (*i.e.*, the failure of glucose levels to increase despite significant reductions in serum insulin levels) reflect the insulin sensitizing, antidiabetic effects of compound 12

25

### Example 20

This example illustrates that compounds of the present invention have potent ability to reduce triglyceride levels in an animal model of type II diabetes.

Male insulin resistant obese Zucker rats were treated by oral administration of compound 12 in a dose of 100 mg/kg/day for one week. The drug was suspended in a mixture of 0.5% carboxymethylcellulose and administered by gavage. Male insulin resistant obese Zucker rats treated with vehicle not containing compound 12

30

represents a useful agent for the prevention and treatment of cardiovascular disease, including the prevention and treatment of restenosis after angioplasty, in diabetics and non-diabetics. Thus, compound 12 has potent ability to reduce triglyceride levels in an animal model of type II diabetes.

5

### Example 21

This example illustrates that compounds of the present invention have ability to inhibit the proliferation of human keratinocytes without inducing cell toxicity.

Human keratinocytes maintained in culture were incubated with varying  
10 concentrations of either compound 12 or troglitazone and the effects on cell proliferation determined by measurement of cell numbers. Compound 12 induced a dose dependent inhibition of cell proliferation in a manner similar to that induced by troglitazone. However, at the highest concentration tested, troglitazone, but not compound 12, was associated with cell toxicity as indicated by increased release of lactate dehydrogenase in  
15 the culture medium when the cell were treated with troglitazone but not compound 12. Thus, compound 12 potently inhibits the proliferation of human keratinocytes without inducing cell toxicity.

### Example 22

20 This example illustrates that compounds of the present invention have ability to inhibit malignant cell proliferation.

Human HT-29 colon cancer cells were incubated with varying concentrations of either compound 12 or troglitazone and the effects on cell proliferation determined by measurement of cell numbers. Compound 12 induced a dose dependent  
25 inhibition of cell proliferation and was more potent than troglitazone in inhibiting malignant cell proliferation. Compound 12 inhibits growth of human colon cancer cells. Human MDA-MB-23 1 breast cancer cells were incubated with varying concentrations of either compound 86 or troglitazone and the effects on cell proliferation determined by measurement of cell numbers. Compound 86 induced a dose dependent inhibition of cell  
30 proliferation and was more potent than troglitazone in inhibiting malignant cell proliferation. Compound 86 inhibits growth of human breast cancer cells. Human MCF-7 breast cancer cells were incubated with varying concentrations of either compound 12 or troglitazone and the effects on cell proliferation determined by measurement of cell

numbers. Compound 12 induced a dose dependent inhibition of malignant cell proliferation. Compound 12 inhibits growth of human breast cancer cells.

### Example 23

5           This example illustrates that compounds of the present invention have ability to inhibit proliferation of 3T3 L 1 fibroblasts and differentiation into adipocytes.

          This example illustrates the effects of compound 87 and troglitazone on proliferation of 3T3 L 1 fibroblasts and differentiation into adipocytes. 3T3L1 cells were incubated with varying concentrations of either compound 87 or troglitazone and the

10       effects on cell proliferation determined by measurement of cell numbers. Compound 12 induced a dose dependent inhibition of fibroblast proliferation in a more potent fashion than troglitazone. However, troglitazone but not compound 87 induced adipogenesis and conversion of the fibroblasts to adipocytes as judged by differential morphology and differential staining of the cells with oil red 0. In vitro molecular assays showed that

15       compound 87 did not activate PPAR gamma. Compound 12 activates PPAR gamma and are potent inducers of adipogenesis. The compounds of the present invention are capable of preventing cell proliferation without inducing adipogenesis and represent useful agents for treatment and prevention of obesity.



**TABLE II:** *Examples of non-malignant proliferative, inflammatory disorders treatable with compounds described in this invention*

<u>Organ System</u>	<u>Disease/Pathology</u>
Dermatological	Psoriasis (all forms), acne vulgaris, acne rosacea, common warts, anogenital (venereal) warts, eczema; lupus associated skin lesions; acute and chronic dermatitides (inflammation of the skin) such as acute and chronic eczema such as atopic dermatitis, allergic dermatitis, contact dermatitis, cosmetic dermatitis, chemical dermatitis, diaper rash, sunburn, seborrheic dermatitis and solar dermatitis; keratoses such as seborrheic keratosis, senile keratosis, actinic keratosis, photo-induced keratosis, skin aging and wrinkle formation including photo-induced skin aging, keratosis follicularis; keloids and prophylaxis against keloid formation; leukoplakia, lichen planus, keratitis, urticaria, pruritus, hidradenitis, acne inversa.
Cardiovascular	Congestive heart failure, endarteritis, endocarditis, atherogenesis, hypertension, vasculo-occlusive diseases including atherosclerosis, thrombosis and restenosis after angioplasty; acute coronary syndromes such as unstable angina, myocardial infarction, ischemic and non-ischemic cardiomyopathies, post-MI cardiomyopathy and myocardial fibrosis, substance-induced cardiomyopathy.
Endocrine	Insulin resistant states including obesity, diabetes mellitus (types 1 & 2), diabetic retinopathy, macular degeneration associated with diabetes, gestational diabetes, impaired glucose tolerance, polycystic ovarian syndrome; osteoporosis, osteopenia, accelerated aging of tissues and organs including Werner's syndrome and Wasting syndrome (all etiologies).
Urogenital	Endometriosis, benign prostatic hypertrophy, leiomyoma, polycystic kidney disease, diabetic nephropathy.
Pulmonary	Asthma, chronic obstructive pulmonary disease (COPD), reactive airway disease, pulmonary fibrosis, pulmonary hypertension.
Immunological/ Connective tissue/ Joints	Rheumatoid arthritis, Raynaud's phenomenon/disease, Sjogren's syndrome systemic sclerosis, systemic lupus erythematosus, vasculitides, ankylosing spondylitis, osteoarthritis, reactive arthritis, psoriatic arthritis, fibromyalgia.
Other	Fibrocystic breast disease, fibroadenoma, chronic fatigue syndrome.

**TABLE IIa: Examples of metabolic, cardiovascular and related proliferative or inflammatory disorders treatable or preventable with compounds described in this invention**

Disorders comprising cardiovascular dysmetabolic syndrome (Syndrome X) and related conditions (prediabetic or diabetic), disease risks and organ complications include:

- 10 • Insulin resistance and/or hyperinsulinemia leading to and including impaired glucose tolerance (prediabetic state) and type 2 (non-insulin dependent) diabetes mellitus.
- Obesity (central obesity) predisposing to the development of insulin resistance.
- 15 • Pancreatic beta-cell dysfunction (e.g. insulinitis secondary to hyperlipemia) and islet cell degeneration occurring later in the natural history of diabetes mellitus.
- Cachexia associated with advanced diabetes mellitus.
- 20 • Hypertension, including insulin resistance-related and/or obesity-related hypertension.
- Dyslipidemia, including abnormally elevated free fatty acids, hypertriglyceridemia, abnormally elevated LDL- and VLDL-cholesterol, conversion of high risk small, dense LDL-cholesterol particles to lower risk, large, buoyant particles),
- 25 hyperapoproteinemia.
- Coagulation abnormalities predisposing to thrombosis, associated with but not limited to: elevated plasminogen activator inhibitor-1, fibrinogen, E-selectin and/or C-reactive protein
- 30 • Type 1 variant of diabetes mellitus, inflammatory and/or T lymphocyte related insulinitis.
- Inflammatory state associated with insulin resistance, mediated by inflammatory cytokines, including but not limited to tumor necrosis factor- $\alpha$  (TNF-  $\alpha$ ) and interleukin-1 (IL-1).
- 35 • Pre-diabetic or diabetic macrovascular complications, including ischemic and occlusive vascular disease (atherosclerosis, atherogenesis), myocardial infarction (MI), stroke, vascular restenosis (arterial or venous) occurring subsequent to invasive procedures.
- 40 • Cardiomyopathy, including post-MI cardiomyopathy.
- 45 • Congestive heart failure (CHF), including CHF related to insulin resistance, obesity and diabetes mellitus (Type 1 and Type 2).
- Endocarditis and endarteritis.

- Pre-diabetic or diabetic microvascular complications, e.g. retinopathy and nephropathy (including prevention of glomerulosclerosis leading to end-stage renal disease).
- Pre-diabetic or diabetic neuropathy related to the central (including autonomic) and/or peripheral nervous system.
- Pre-diabetic or diabetic ocular diseases, including diabetic retinopathy, retinal ischemia, cataract, macular degeneration.
- Insulin resistance, hypertension, dyslipidemia and cardiovascular risk factors associated with polycystic ovarian syndrome

**TABLE III: Examples of neoplastic diseases or malignancies diseases treatable with compounds described in this invention**

Organ System	Malignancy/Cancer type
Skin	Basal cell carcinoma, melanoma, squamous cell carcinoma; cutaneous T cell lymphoma; Kaposi's sarcoma.
Hematological	Acute leukemia, chronic leukemia and myelodysplastic syndromes.
Urogenital	Prostatic, renal and bladder carcinomas, anogenital carcinomas including cervical, ovarian, uterine, vulvar, vaginal, and those associated with human papilloma virus infection.
Neurological	Gliomas including glioblastomas, astrocytoma, ependymoma, medulloblastoma, oligodendroma; meningioma, pituitary adenoma, neuroblastoma, craniopharyngioma.
Gastrointestinal	Colon, colorectal, gastric, esophageal, mucocutaneous carcinomas.
Breast	Breast cancer including estrogen receptor and progesterone receptor positive or negative subtypes, soft tissue tumors.
Metastasis	Metastases resulting from the neoplasms.
Other	Angiomata, angiogenesis associated with the neoplasms.

**TABLE IIIa:**

Location	Malignancy/Cancer type
Various	fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilms' tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, melanoma, neuroblastoma, and retinoblastoma.

**TABLE IV:** *Examples of viral infections and related pathologies treatable with compounds described in this invention*

Virus	Viral infection/cancer or other virus-associated pathology
HTLV	T-cell leukemia/lymphoma, HTLV-associated arthritides/myelopathies.
HPV	Cervical and anogenital cancers; common and anogenital (venereal) warts, including verrucae, condyloma or condyloma acuminata, related non-neoplastic (e.g., keratitis, conjunctivitis) pre-neoplastic and neoplastic (e.g., conjunctival epithelial neoplasms) diseases of the eye.
HAV,HBV,HCV	Hepatitis, hepatocellular carcinoma, lymphoma.
CMV	Hepatitis, retinitis, meningitis.
HSV, VSV	Related mucocutaneous, oropharyngeal and genital diseases, related skin and respiratory infections, varicella-zoster, chicken pox, herpes zoster, post-herpetic neuralgia, conjunctivitis, keratoconjunctivitis, keratitis.
HHV	Exanthem subitum, infectious mononucleosis.
EBV	Infectious mononucleosis, chronic fatigue syndrome, lymphoma, conjunctivitis, keratitis, and related infections of the eye.
Adenoviruses	Upper and lower respiratory tract infections, pneumonia, conjunctivitis.
RSV	Upper and lower respiratory tract infections, pneumonia.
PMV	Mumps and related manifestations, e.g., conjunctivitis.
MV, RV	Measles, Rubella ("German measles") and related manifestations.
Coxsackie viruses	Conjunctivitis, diabetes mellitus, respiratory infections.
Influenza viruses	Upper and lower respiratory tract infections, pneumonia.
HIV, Human Immunodeficiency Virus; HTLV, Human T-cell Lymphocyte Virus; HPV, Human Papilloma Virus; HAV, Hepatitis A Virus; HBV, Hepatitis B Virus; HAV, Hepatitis C Virus; CMV, Cytomegalovirus; HSV, Herpes Simplex Virus (Types I & II); HHV, Human Herpes Virus; EBV, Epstein-Barr Virus; RSV, Respiratory Syncytial Virus; VZV, Varicella-Zoster Virus; PMV, Paramyxovirus; MV, Measles (Rubeola) Virus; RV, Rubella Virus	

**Table V:** *HIV related infections and diseases treatable with compounds described in this invention*

Organ system	Viral infection/manifestation or other HIV-associated disease
Immunologic	AIDS, primary HIV infection.
Dermatological	Anogenital cancers including rectal and cervical cancer, Kaposi's sarcoma, atopic dermatitis, squamous cell carcinoma, hairy leukoplakia, molluscum contagiosum, warts (HPV infections), seborrheic dermatitis, psoriasis, xeroderma, HSV and varicella-zoster infections.
Hematologic	Non-Hodgkin's lymphoma, B cell lymphoma, anemia, neutropenia, thrombocytopenia.
Gastrointestinal	Anorexia, gastroparesis, diarrhea, malabsorption, gastrointestinal CMV infections, esophagitis, colitis, hepatitis, lymphoma.
Ocular	Conjunctivitis, keratitis, keratoconjunctivitis, uveitis, retinitis, chorioretinitis, CMV retinitis, iridocyclitis, vitreitis, choroiditis, papilledema, Kaposi's sarcoma, lymphoma, ocular palsies, conjunctival warts, pre-neoplastic and neoplastic diseases of the eye.
CardiacMyocarditis,	endocarditis, pericarditis.
Pulmonary	CMV pneumonitis, lymphoid interstitial pneumonitis.
Nephrologic	HIV nephropathy, renal cell carcinoma, amyloidosis, uropathy.
Rheumatologic	Arthralgia, fibromyalgia, Reiter's syndrome, psoriatic arthritis, vasculitis.
Neurologic	Dementia, viral meningitis, viral encephalitis, HIV encephalopathy, progressive multifocal leukoencephalopathy, CNS lymphoma, peripheral and autonomic neuropathies.
Psychiatric	Dysphoric mood disorders, depression, depression associated with chronic diseases and medications, bipolar disorder, anxiety disorders, chronic fatigue syndrome, chronic pain, psychoses, substance abuse disorders and drug addiction.
General	Lymphoma, metastatic lymphoma, Kaposi's sarcoma, wasting syndrome.

**TABLE VI: Examples of neurological and psychiatric disorders that can be treated with compounds described in this invention**

---

5	<u>Neurological Disorders</u>
	Migraine headaches
	Pain disorders including algesia, hyperalgesia, acute and chronic pain, allodynia
	Chronic fatigue syndrome
	Amnesia
10	<u>Psychiatric Disorders</u>
	Dysphoric mood disorders
	Dysthymic disorder
	Depression including depression associated with chronic diseases and medications
15	Manic depressive disorder
	Anxiety states including panic disorder and agoraphobia
	Post menstrual syndrome
	Obsessive-compulsive disorder, schizophrenia, chronic fatigue syndrome
	Substance abuse and drug addiction
20	Anorexia nervosa
	Bulimia nervosa
	<u>Neurodegenerative Disorders</u>
	Alzheimer's disease and other dementias
25	Parkinson's disease
	Huntington's chorea
	Upper and lower motor neuron degenerative diseases
	amyotrophic lateral sclerosis
	primary lateral sclerosis
30	spinal muscular atrophy and dystrophy
	Multiple sclerosis and other demyelinating diseases
	Progressive ataxias
	Post-anoxia ataxias and CNS degeneration
	Drug and ethanol-induced ataxias and CNS degeneration
35	<u>Neurological Disorders</u>
	Migraine headaches
	Pain disorders including algesia, hyperalgesia, acute and chronic pain, allodynia
	Chronic fatigue syndrome
40	Amnesia

---

**TABLE VIa: *Prevention of allograft rejection and amelioration of the associated syndrome complex***

---

- 5 The compounds of the present invention are useful in organ transplantation, especially  
uses pertaining to the prevention and treatment of the syndrome complex encountered in  
allograft rejection, and in kidney, heart, liver and skin transplantation, and uses pertaining  
to the promotion of allograft survival. This includes inflammatory and proliferative  
conditions or diseases associated with allograft transplantation and immune suppression  
10 including: acute allograft rejection, chronic allograft rejection, graft versus host disease,  
post-transplantation de novo malignancies (e.g. lymphoma and epidermal cancers),  
osteoporosis and osteopenia, hyperlipidemia, insulin resistance and diabetes mellitus,  
hypertension, atherosclerosis, endarteritis associated with heart allograft transplantation,  
glomerulonephritis associated with renal allograft transplantation, cardiomyopathy and  
15 congestive heart failure associated with allograft transplantation, and in particular, heart  
transplantation.



**TABLE VII: Diseases of the eye treatable with compounds described in this invention****1. Diseases caused by viruses or associated with viral infections**

<b>Disease</b>	<b>Virus</b>
Blepharitis	HSV, VZV, Vaccinia, HPV, molluscum contagiosum
Conjunctivitis	HSV, VZV, EBV, Adenovirus, Vaccinia, Variola, HPV, molluscum contagiosum, influenza
Follicular c.	Newcastle, measles, mumps, rubella, molluscum contagiosum
Hemorrhagic c.	Enterovirus, coxsackie
Catarrhal c	Rubella
Keratitis	HSV, VZV, EBV, Adenovirus, Vaccinia, Variola, HPV, molluscum contagiosum
Keratoconjunctivitis	HSV, VZV, EBV, Adenovirus, Vaccinia, Variola, HPV, molluscum contagiosum
Retinitis	CMV
Uveitis	HPV
Conjunctival warts	HPV
C. epithelial neoplasms	HPV

**2. Ocularplastic diseases**

**Benign tumors:** Keratocanthoma, molluscum contagiosum, dermoid cysts, neurofibroma, neurofibromatosis, schwannoma (neurilemoma), pleiomorphic adenoma

**Malignant tumors:** Basal cell carcinoma, squamous cell carcinoma, mucoepidermoid carcinoma, melanoma, retinoblastoma, embryonal rhabdomyosarcoma, meningioma, adenoid cystic carcinoma, lymphoid tumors of the orbit, mesenchymal tumors (fibrous hystiocytoma) of the orbit, nasopharyngeal carcinoma.

**Vascular lesions:** Hemangioma, lymphangioma.

**3. Inflammatory/immunological ocular diseases**

Acute allergic conjunctivitis and hypersensitivity reactions  
 Drug-related inflammation and hypersensitivity reactions  
 Chronic (vernal) conjunctivitis  
 Contact lens-associated conjunctivitis, e.g. giant papillary conjunctivitis  
 Conjunctival ulceration, including ulceration associated with mucous membrane pemphigoid and the Steven's-Johnson syndrome, leading to progressive fibrosis and scarring, cicatrization and symblepharon.  
 Corneal abrasion, corneal ulceration, or corneal injury of any etiology.

**4. Other lesions**


Retina: Macular degeneration, retinopathy, including diabetic r. and hypertensive r.  
 Lens: Cataract, all etiologies including rheumatological and collagen vascular diseases  
 Uvea: Ueitis, vitreitis, all etiologies, including UV radiation and diabetes.

All publications, patents and patent applications mentioned in this specification are herein incorporated by reference into the specification in their entirety for all purposes.

5           Although the invention has been described with reference to preferred  
embodiments and examples thereof, the scope of the present invention is not limited only  
to those described embodiments. As will be apparent to persons skilled in the art,  
modifications and adaptations to the above-described invention can be made without  
departing from the spirit and scope of the invention, which is defined and circumscribed  
10 by the appended claims.

**WHAT IS CLAIMED IS:**

1. A compound of the formula



wherein:

$R^1$  and  $R^2$  are each independently a member selected from the group consisting of hydrogen,  $C(O)-R^6$  and  $C(S)-R^6$ , wherein  $R^6$  is a member selected from the group consisting of hydrogen,  $(C_1-C_{12})$ alkyl, aryl, arylalkyl,  $(C_1-C_{12})$ carboxyl,  $(C_1-C_{12})NHR^7$ ,  $(C_1-C_{12})NR^7R^8$ ,  $OR^7$ ,  $NHR^7$ ,  $SR^7$  and  $NR^7R^8$ , wherein  $R^7$  and  $R^8$  are each independently a member selected from the group consisting of hydrogen,  $(C_1-C_{12})$ alkyl, aryl and arylalkyl; or

$R^1$  and  $R^2$  together with the sulfurs to which they are bound join to form a 1,2-dithiolane ring;

X is a member selected from the group consisting of O, NR,  $C(O)O$ ,  $OC(O)O$  and  $C(O)NR$ , wherein R is a member selected from the group consisting of hydrogen, optionally substituted  $(C_1-C_6)$ alkyl and optionally substituted aryl;

Y is a member selected from the group consisting of O, S and  $NR^3$ , wherein  $R^3$  is a member selected from the group consisting of hydrogen and optionally substituted  $(C_1-C_6)$ alkyl;


n is an integer from 2 to 14;

m is an integer from 0 to 14;

q is an integer from 0 to 1; and

t is an integer from 0 to 1, or a pharmaceutical acceptable salt or solvate thereof, with the proviso that when m is 0 then q is 0.

2. A compound in accordance with claim 1, having the formula



wherein:

$R^1$  and  $R^2$  are each independently a member selected from the group consisting of hydrogen,  $C(O)-R^6$  and  $C(S)-R^6$ , wherein  $R^6$  is a member selected from the group consisting hydrogen,  $(C_1-C_{12})$ alkyl, aryl, arylalkyl,  $(C_1-C_{12})$ carboxyl,  $(C_1-C_{12})NHR^7$ ,  $(C_1-C_{12})NR^7R^8$ ,  $OR^7$ ,  $NHR^7$ ,  $SR^7$  and  $NR^7R^8$ , wherein  $R^7$  and  $R^8$  are each independently a member selected from the group consisting of hydrogen,  $(C_1-C_{12})$ alkyl, aryl and arylalkyl; or

$R^1$  and  $R^2$  together with the sulfurs to which they are bound join to form a 1,2-dithiolane ring;

X is a member selected from the group consisting of O, NR,  $C(O)O$ ,  $OC(O)O$  and  $C(O)NR$ , wherein R is a member selected from the group consisting of hydrogen, optionally substituted  $(C_1-C_6)$ alkyl and optionally substituted aryl;

Y is a member selected from the group consisting of O, S and  $NR^3$ , wherein  $R^3$  is a member selected from the group consisting of hydrogen and optionally substituted  $(C_1-C_6)$ alkyl;

n is an integer from 2 to 14;

m is an integer from 0 to 14;

q is an integer from 0 to 1; and

t is an integer from 0 to 1 with the proviso that when m is 0 then q is 0.

3. A compound in accordance with claim 2, wherein:

m is an integer from 1 to 6;

q is 1; and

t is 1.

4. A compound in accordance with claim 3, wherein:

$R^1$  and  $R^2$  join to form a 1,2-dithiolane ring;

X is  $C(O)NR$ , wherein R is a member selected from the group consisting of hydrogen and optionally substituted  $(C_1-C_6)$ alkyl;

5 Y is O;  
6 n is 4; and  
7 m is 2.

1 5. A compound in accordance with claim 4, wherein:  
2 R is methyl.

1 6. A compound in accordance with claim 3, wherein:  
2 R<sup>1</sup> is hydrogen;  
3 R<sup>2</sup> is hydrogen;  
4 X is C(O)NR, wherein R is a member selected from the group consisting  
5 of hydrogen and optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl;  
6 y is O;  
7 n is 4; and  
8 m is 2.

1 7. A compound in accordance with claim 6, wherein:  
2 R is methyl.

1 8. A compound in accordance with claim 3, wherein:  
2 R<sup>1</sup> and R<sup>2</sup> join to form a 1,2-dithiolane ring;  
3 X is C(O)O  
4 y is O;  
5 n is 4; and  
6 m is 2.

1 9. A compound in accordance with claim 3, wherein:  
2 R<sup>1</sup> is hydrogen;  
3 R<sup>2</sup> is hydrogen;  
4 X is C(O)O;  
5 y is O;  
6 n is 4; and  
7 m is 2.

1 10. A compound in accordance with claim 3, wherein:

2             $R^1$  and  $R^2$  join to form a 1,2-dithiolane ring;  
3            x is O;  
4            y is O;  
5            n is 5; and  
6            m is 2.

1            11.    A compound in accordance with claim 3, wherein:  
2             $R^1$  is hydrogen;  
3             $R^2$  is hydrogen;  
4            x is O;  
5            y is O;  
6            n is 5; and  
7            m is 2.

1            12.    A compound in accordance with claim 3, wherein:  
2             $R^1$  and  $R^2$  join to form a 1,2-dithiolane ring;  
3            X is OC(O)O;  
4            Y is O;  
5            n is 4; and  
6            m is 2.

1            13.    A compound in accordance with claim 2, wherein:  
2             $R^1$  is hydrogen;  
3             $R^2$  is hydrogen;  
4            X is OC(O)O;  
5            Y is O;  
6            n is 4; and  
7            m is 2.

1            14.    A compound in accordance with claim 2, wherein:  
2            m is 0;  
3            q is 0; and  
4            t is 1.

1            15.    A compound in accordance with claim 14, wherein:

2           R<sup>1</sup> and R<sup>2</sup> join to form a 1,2 dithiolane ring;  
3           X is C(O)NR, wherein R is a member selected from the group consisting  
4 of hydrogen and optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl; and  
5           n is 4.

1           **16.**   A compound in accordance with claim 15, wherein:  
2           R is methyl.

1           **17.**   A compound in accordance with claim 14, wherein:  
2           R<sup>1</sup> is hydrogen;  
3           R<sup>2</sup> is hydrogen;  
4           X is C(O)NR, wherein R is a member selected from the group consisting  
5 of hydrogen and optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl; and  
6           n is 4.

1           **18.**   A compound in accordance with claim 17, wherein:  
2           R is methyl.

1           **19.**   A compound in accordance with claim 14, wherein:  
2           R<sup>1</sup> and R<sup>2</sup> join to form a 1,2-dithiolane ring;  
3           X is O; and  
4           n is 5.

1           **20.**   A compound in accordance with claim 14, wherein:  
2           R<sup>1</sup> is hydrogen;  
3           R<sup>2</sup> is hydrogen;  
4           X is O; and  
5           n is 5.

1           **21.**   A compound in accordance with claim 2, wherein:  
2           m is an integer from 2 to 14;  
3           q is 0; and  
4           t is 0.

1           **22.**   A compound in accordance with claim 21, wherein:

2                   R<sup>1</sup> and R<sup>2</sup> join to form a 1,2-dithiolane ring;  
 3                   X is C(O)NR, wherein R is a member selected from the group consisting  
 4 of hydrogen and optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl;  
 5                   n is 4; and  
 6                   m is 3.

1                   **23.**    A compound in accordance with claim 22, wherein:  
 2                   R is methyl.

1                   **24.**    A compound in accordance with claim 21, wherein:  
 2                   R<sup>1</sup> and R<sup>2</sup> join to form a 1,2 dithiolane ring;  
 3                   X is O;  
 4                   n is 5; and  
 5                   m is 3.


1                   **25.**    A compound in accordance with claim 21, wherein:  
 2                   R<sup>1</sup> is hydrogen;  
 3                   R<sup>2</sup> is hydrogen;  
 4                   X is O;  
 5                   n is 5; and  
 6                   m is 3.

1                   **26.**    A compound in accordance with claim 2, wherein:  
 2                   said compound is a member selected from the group consisting of N-(2-{4-  
 3 [(2,4-dioxo(1,3-thiazolidin-5-yl)methyl]phenoxy}ethyl)-5-(1,2-dithiolan-3-yl)  
 4 pentanamide, N-(2-{4-[(2,4-dioxo(1,3-thiazolidin-5-yl)methyl]phenoxy}ethyl)-5-(1,2-  
 5 dithiolan-3-yl)-N-methylpentanamide, N-{4-[(2,4-dioxo(1,3-thiazolidin-5-  
 6 yl)methyl]phenyl}-5-(1,2-dithiolan-3-yl)-N-methyl pentanamide, N-(2-{4-[(2,4-  
 7 dioxo(1,3-thiazolidin-5-yl)methyl]phenoxy}ethyl)-5-(1,2-dithiolan-3-yl)-N-  
 8 methylacetamide, 5-[(4-{2-[5-(1,2-dithiolan-3-  
 9 yl)pentyl)methylamino]ethoxy}phenyl)methyl]-1,3-thiazolidine-2,4-dione, tartrate salt of  
 10 5-[(4-{2-[5-(1,2-dithiolan-3-yl)pentyl)methylamino]ethoxy}phenyl) methyl]-1,3-  
 11 thiazolidine-2,4-dione, N-(2-{4-[(2,4-dioxo(1,3-thiazolidin-5-yl)methyl]phenoxy}ethyl)-



- 12 N-methyl-6,8-disulfanyloctamide and 6,8-diacetylthio-N-(2-{4-[(2,4-dioxo(1,3-  
13 thiazolidin-5-yl)methyl]phenoxy}ethyl)-N-methyloctanamide.

1 27. A pharmaceutical composition comprising a compound of the  
2 formula compound of the formula



3  
4 wherein:

5 R<sup>1</sup> and R<sup>2</sup> are each independently a member selected from the group  
6 consisting of hydrogen, C(O)-R<sup>6</sup> and C(S)-R<sup>6</sup>, wherein R<sup>6</sup> is a member selected from the  
7 group consisting of hydrogen, (C<sub>1</sub>-C<sub>12</sub>)alkyl, aryl, arylalkyl, (C<sub>1</sub>-C<sub>12</sub>)carboxyl, (C<sub>1</sub>-  
8 C<sub>12</sub>)NHR<sup>7</sup>, (C<sub>1</sub>-C<sub>12</sub>)NR<sup>7</sup>R<sup>8</sup>, OR<sup>7</sup>, NHR<sup>7</sup>, SR<sup>7</sup> and NR<sup>7</sup>R<sup>8</sup>, wherein R<sup>7</sup> and R<sup>8</sup> are each  
9 independently a member selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>12</sub>)alkyl,  
10 aryl and arylalkyl; or

11 R<sup>1</sup> and R<sup>2</sup> together with the sulfurs to which they are bound join to form a  
12 1,2-dithiolane ring;

13 X is a member selected from the group consisting of O, NR, C(O)O,  
14 OC(O)O and C(O)NR, wherein R is a member selected from the group consisting of  
15 hydrogen, optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl and optionally substituted aryl;

16 Y is a member selected from the group consisting of O, S and NR<sup>3</sup>,  
17 wherein R<sup>3</sup> is a member selected from the group consisting of hydrogen and optionally  
18 substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl;

19 n is an integer from 2 to 14;

20 m is an integer from 0 to 14;

21 q is an integer from 0 to 1;

22 t is an integer from 0 to 1, or a pharmaceutical acceptable salt or solvate  
23 thereof; with the proviso that when m is 0 then q is 0; and

24 a pharmaceutical acceptable carrier.

1 28. A composition in accordance with claim 27, wherein:

2 M is an integer from 1 to 6;

3 q is 1; and

4 t is 1.

1 29. A composition in accordance with claim 27, wherein:

2 m is 0;

3 q is 0; and

4 t is 1.


1 30. A composition in accordance with claim 27, wherein:

2 m is an integer from 2 to 14;

3 q is 0; and

4 t is 0.

1 31. A method of treating a PPAR $\gamma$  mediated disease or oxidative stress,  
2 said method comprising administering to a subject a therapeutically effective amount of a  
3 compound of the of the formula



4  
5 wherein:

6 R<sup>1</sup> and R<sup>2</sup> are each independently a member selected from the group  
7 consisting of hydrogen, C(O)-R<sup>6</sup> and C(S)-R<sup>6</sup>, wherein R<sup>6</sup> is a member selected from the  
8 group consisting of hydrogen, (C<sub>1</sub>-C<sub>12</sub>)alkyl, aryl, arylalkyl, (C<sub>1</sub>-C<sub>12</sub>)carboxyl, (C<sub>1</sub>-  
9 C<sub>12</sub>)NHR<sup>7</sup>, (C<sub>1</sub>-C<sub>12</sub>)NR<sup>7</sup>R<sup>8</sup>, OR<sup>7</sup>, NHR<sup>7</sup>, SR<sup>7</sup> and NR<sup>7</sup>R<sup>8</sup>, wherein R<sup>7</sup> and R<sup>8</sup> are each  
10 independently a member selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>12</sub>)alkyl,  
11 aryl and arylalkyl; or

12 R<sup>1</sup> and R<sup>2</sup> together with the sulfurs to which they are bound join to form a  
13 1,2-dithiolane ring;

14 X is a member selected from the group consisting of O, NR, C(O)O,  
15 OC(O)O and C(O)NR, wherein R is a member selected from the group consisting of  
16 hydrogen, optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl and optionally substituted aryl;


17 Y is a member selected from the group consisting of O, S and NR<sup>3</sup>,  
 18 wherein R<sup>3</sup> is a member selected from the group consisting of hydrogen and optionally  
 19 substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl;  
 20 n is an integer from 2 to 14;  
 21 m is an integer from 0 to 14;  
 22 q is an integer from 0 to 1; and  
 23 t is an integer from 0 to 1; or a pharmaceutical acceptable salt or solvate  
 24 thereof; with the proviso that when m is 0 then q is 0.

1 32. A method in accordance with claim 31, wherein:  
 2 m is an integer from 1 to 6;  
 3 q is 1; and  
 4 t is 1.

1 33. A method in accordance with claim 31, wherein:  
 2 m is 0;  
 3 q is 0; and  
 4 t is 1.

1 34. A method in accordance with claim 31, wherein:  
 2 m is an integer from 2 to 14;  
 3 q is 0; and t is 0.

1 35. A compound of the formula



2  
 3 wherein:

4 R<sup>1</sup> and R<sup>2</sup> are each independently a member selected from the group  
 5 consisting of hydrogen, C(O)-R<sup>6</sup> and C(S)-R<sup>6</sup>, wherein R<sup>6</sup> is a member selected from the  
 6 group consisting of hydrogen, (C<sub>1</sub>-C<sub>12</sub>)alkyl, aryl, arylalkyl, (C<sub>1</sub>-C<sub>12</sub>)carboxyl, (C<sub>1</sub>-  
 7 C<sub>12</sub>)NHR<sup>7</sup>, (C<sub>1</sub>-C<sub>12</sub>)NR<sup>7</sup>R<sup>8</sup>, OR<sup>7</sup>, NHR<sup>7</sup>, SR<sup>7</sup> and NR<sup>7</sup>R<sup>8</sup>, wherein R<sup>7</sup> and R<sup>8</sup> are each

8 independently a member selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>12</sub>)alkyl,  
9 aryl and arylalkyl; or

10 R<sup>1</sup> and R<sup>2</sup> together with the sulfurs to which they are bound join to form a  
11 1,2-dithiolane ring;

12 X is a member selected from the group consisting of O, NR, C(O)O,  
13 OC(O)O and C(O)NR, wherein R is a member selected from the group consisting of  
14 hydrogen, optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl and optionally substituted aryl;

15 Y is a member selected from the group consisting of O, S and NR<sup>3</sup>,  
16 wherein R<sup>3</sup> is a member selected from the group consisting of hydrogen and optionally  
17 substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl;

18 n is an integer from 2 to 14;

19 m is an integer from 0 to 14;

20 q is an integer from 0 to 1; and

21 t is an integer from 0 to 1 with the proviso that when m is 0 then q is 0.

22 R<sup>5</sup> is a member selected from the group consisting of hydrogen and  
23 optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl;

24 n is an integer from 2 to 14;

25 m is an integer from 0 to 14; and

26 q is an integer from 0 to 1 or a pharmaceutical acceptable salt or solvate  
27 thereof; with the proviso that when m is 0 then q is 0.

Figure 1






Figure 2

A



B



C




Figure 3




Figure 4






Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




Figure 12






Figure 13











Figure 14



Tartrate Salt

Figure 15



## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US00/05895**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(7) : Please See Extra Sheet.

US CL : 548/183; 549/39; 560/9; 562/426 514/369, 440,538,562

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 548/183; 549/39; 560/9; 562/426 514/369, 440,538,562

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JP 08-201957 A (YAMADA) 09 August 1996 entire document.	1



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*B* earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A*	document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means		
*P* document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

28 JUNE 2000

Date of mailing of the international search report

11 JUL 2000

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

ROBERT GERSTL

Telephone No. (703) 308-1235

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/05895

## A. CLASSIFICATION OF SUBJECT MATTER:

IPC (7):

C07D 417/12, 339/04; C07C 327/18, 327/36; A61K 31/216, 31/192, 31/427, 31/385